Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 5 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Impact of Water Management and Soil Properties on Rice Productivity in Paddy Fields of Punjab

Waseem Hassan
Department of Soil and Environmental Sciences, Muhammad Nawaz Shareef University of Agriculture Multab Email: waseem.hassan@mnsuam.edu.pk

Muhammad Aaqib
Department of Agronomy, University of Agriculture, Faisalabad
Email: muhammadaaqibuaf@gmail.com

Mahdia
MSc Soil Science, Institute of soil and environmental Sciences, University of Agriculture Faisalabad Email: mahdiakhalid786@gmail.com

Abdul Khaliq
Soil And Water Testing Laboratory Kotla Ahmad Road Rajanpur
Email: gishkori757@gmail.com

Fareeha Habib*
Senior Scientist (laboratory), Soil and Water Testing Laboratory for Research, Dera Ghazi Khan Email: fareehahabib.ao@ gmail.com

Muhammad Aslam
Senior Scientist Pesticide Quality Control Lab Multan
Email: maslamswtldgk85@gmail.com

Marghoob Ahmad
Centre of Agricultural Biochemistry and Biotechnology (CABB), University of Agriculture Faisalabad, Pakistan Email: marghoobahmad1219@gmail.com

ABSTRACT

Rice (Oryza sativa L.) is the most strategically important cereal crop of South Asia, occupying a central position in the food systems of Pakistan and India. The Punjab province, shared between both nations and forming the agricultural heartland of the region, contributes the largest share of rice production in the subcontinent. However, mounting challenges related to water scarcity, soil degradation, and climate variability now threaten the long term productivity of paddy systems in this region. This review synthesizes the existing body of literature on how water management practices and soil physicochemical properties jointly influence rice yield, water use efficiency, and ecosystem sustainability in Punjab paddy fields. Key themes addressed include the hydrological dynamics of flooded and non flooded paddy cultivation, the agronomic

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 5 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

performance of irrigation scheduling techniques such as alternate wetting and drying (AWD) and saturated soil culture (SSC), the impact of soil organic matter, soil texture, and nutrient cycling on rice productivity, and the interactive effects of waterlogging on soil redox chemistry and nitrogen transformations. Evidence from field trials, remote sensing analyses, and modeling studies across the Indo Gangetic Plain is reviewed and synthesized. The paper also addresses methane emissions and broader environmental trade offs associated with different water management regimes. Recommendations are provided for research gaps and future directions in sustainable paddy agronomy in Punjab.

Keywords: Rice Productivity, Water Management, Soil Properties, Punjab, Alternate Wetting And Drying, Paddy Agronomy, Nitrogen Cycling, Waterlogging

Introduction

Rice is a dietary staple for more than half of the global human population and serves as the primary caloric source for over 3.5 billion people across Asia, Sub Saharan Africa, and Latin America (Khush, 2005). South Asian nations, particularly Pakistan and India, are among the world's foremost rice producers, and the Punjab province of both countries together accounts for approximately 25 to 30 percent of the combined national rice output (Pakistan Bureau of Statistics, 2022; Ministry of Agriculture India, 2023). The ecological and agronomic conditions of Punjab, including its fertile alluvial soils derived from the five major rivers of the Indus basin, a sub tropical climate with pronounced monsoon rainfall, and a well developed canal irrigation network, have historically supported high rice yields. However, current trajectories of groundwater depletion, soil salinity intrusion, and climate driven shifts in precipitation patterns raise serious concerns about the sustainability of existing paddy cultivation systems (Humphreys et al., 2010).

Water management in paddy fields is uniquely complex compared to other cereal crops. Traditional practice involves continuous flooding from transplanting until shortly before harvest, a regime that satisfies the anaerobic physiology of transplanted rice but simultaneously consumes enormous volumes of irrigation water, often exceeding 1,200 to 1,500 millimeters per crop season (Bouman and Tuong, 2001). In Punjab, where canal water availability is increasingly unreliable and groundwater tables in districts such as Sheikhupura, Hafizabad, and Gujranwala are declining at rates between 0.3 and 1.0 meters per year (Shah et al., 2009), the profligate use of water in paddy cultivation is becoming economically and ecologically untenable. Simultaneously, the soil resource base of Punjab paddy fields, while inherently fertile, is under threat from long term monoculture, excessive tillage, organic matter decline, and secondary salinization, all of which erode the physical and chemical properties that underpin productive rice growth (Timsina and Connor, 2001).

This review critically examines the scientific literature addressing the intersection of water management strategies and soil property dynamics as they relate to rice productivity in Punjab paddy systems. The objective is threefold: first, to assess the agronomic and hydrological evidence for water saving irrigation technologies; second, to evaluate how key soil properties including texture, organic carbon, pH, bulk density, and nutrient pools respond to different water management regimes; and

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 5 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

third, to identify opportunities for integrating soil and water management in ways that simultaneously sustain or increase rice yields while conserving natural resources.

Rice Cultivation in Punjab: Regional Context and Agronomic Significance

Geographic and Climatic Overview

Punjab occupies a vast alluvial plain stretching across both Pakistani and Indian territory, bounded by the Himalayas to the north, the Thar Desert to the south, and intersected by the tributaries of the Indus river system. Mean annual rainfall ranges from 250 to 500 millimeters in Pakistani Punjab to 400 to 800 millimeters in Indian Punjab, supplemented extensively by canal and groundwater irrigation (Arshad et al., 2008). The rice growing season in Punjab corresponds broadly to the Kharif cropping period, extending from June or July transplanting through October or November harvest, aligning with peak monsoon moisture. Mean temperatures during the growing season range from 28 to 35 degrees Celsius, conditions that are broadly favorable for Indica type rice varieties dominant in the region.

Soil types across the Punjab rice belt are predominantly loamy to clay loam in texture, formed from Holocene alluvial deposits of variable mineralogy. These soils are generally fertile, with moderate to high cation exchange capacity and a natural tendency toward neutral to slightly alkaline pH, though waterlogging during the Kharif season temporarily induces reducing conditions that alter nutrient dynamics profoundly (Sahrawat, 2004).

Rice Varieties and Cultivation Systems

The dominant rice cultivation system in Punjab is the transplanted rice monoculture or the rice wheat rotation, in which paddy rice follows or precedes wheat (Triticum aestivum L.) in an annual double crop cycle. This rotation occupies an estimated 10 to 12 million hectares across the two Punjab provinces and forms the backbone of regional food security (Ladha et al., 2003). Commonly cultivated rice varieties include Basmati 385, Super Basmati, Basmati 515, and Kainat in Pakistani Punjab, while IR series varieties and Parmal types are more common in irrigated non Basmati areas of Indian Punjab. Yield potential varies considerably between varieties and cropping systems, with Basmati types typically yielding 3 to 4.5 tonnes per hectare under optimal conditions and high yielding coarse rice varieties reaching 5 to 7 tonnes per hectare (IRRI, 2023).

Transplanting is the predominant method of rice establishment in Punjab, with 25 to 30 day old seedlings transferred to puddled fields in June or July. Puddling, the wet tillage of saturated soils to disrupt soil structure and reduce percolation losses, is a standard practice that fundamentally alters soil physical properties and has both agronomic benefits and long term soil health implications (Sharma and De Datta, 1985).

Water Management in Punjab Paddy Systems

Conventional Continuous Flooding

Continuous flooding (CF) has been the default water management practice for transplanted paddy rice in South Asia for millennia. Under CF, fields are maintained under 5 to 10 centimeters of standing water from shortly after transplanting until

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 5 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

approximately 10 to 14 days before harvest. This regime creates the anaerobic rhizosphere conditions that rice tolerates through aerenchyma mediated oxygen transport and that suppress many weed species competing with rice (Neue and Lantin, 1994). However, CF is an extraordinarily water intensive system. Field measurements across the Indo Gangetic Plain consistently report seasonal water inputs under CF ranging from 900 to 2,100 millimeters, of which only 450 to 650 millimeters is consumed as evapotranspiration (ET), meaning that 40 to 70 percent of applied water is lost to unproductive pathways including deep percolation, seepage, and surface runoff (Tuong and Bouman, 2003).

In the context of Punjab's groundwater crisis, these losses are both economically damaging and environmentally unsustainable. The energy cost of groundwater pumping for paddy irrigation in Indian Punjab alone is estimated to exceed 15 billion rupees annually, and Pakistani Punjab faces similarly escalating pump irrigation costs as aquifer levels decline (Mukherji, 2007). The environmental consequences of CF also include significant emissions of methane (CH4), a potent greenhouse gas produced by methanogenic archaea under anoxic paddy soil conditions, with global paddy rice estimated to contribute 25 to 100 teragrams of CH4 annually (Linquist et al., 2012).

Alternate Wetting and Drying

Alternate wetting and drying (AWD) has emerged over the past two decades as the most extensively researched and field validated water saving technology for irrigated paddy rice. Under AWD, fields are allowed to dry following each flooding event until the water table drops to a defined threshold below the soil surface, typically 15 to 20 centimeters below the soil surface for safe AWD or as little as 5 centimeters for moderate AWD, before re irrigation is applied (Bouman et al., 2007). AWD can be implemented at different intensities depending on the desired balance between water savings and potential yield risk.

Field trials conducted across Pakistan and India have consistently demonstrated that AWD reduces seasonal irrigation water input by 15 to 30 percent compared to CF without statistically significant reductions in grain yield when the water table threshold during drying is maintained above 15 to 20 centimeters below the soil surface (Belder et al., 2004; Lampayan et al., 2015). A meta analysis by Carrijo et al. (2017) encompassing 53 studies found that moderate AWD reduced CH4 emissions by 48 percent and maintained yields within 5 percent of CF in well managed systems. In Pakistani Punjab specifically, trials at the Pakistan Rice Research Institute (PRRI) in Kala Shah Kaku demonstrated water savings of 20 to 25 percent under AWD with yield equivalence to CF for Super Basmati rice (Tabbal et al., 2002).

The agronomic rationale for AWD lies in rice's ability to maintain root water uptake during moderate soil drying through a combination of root hydraulic redistribution and stomatal regulation, while the aerobic intervals during drying periods promote soil nitrogen mineralization and reduce ferrous iron toxicity in reduced soils (Peng et al., 2004). However, AWD implementation in Punjab faces practical barriers including lack of farmer awareness, absence of reliable field water tubes for monitoring soil water status, and risk aversion among smallholders who fear yield losses from over drying during critical growth stages such as panicle initiation and

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 5 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

grain filling.

Saturated Soil Culture

Saturated soil culture (SSC) is a more conservative water saving approach in which the paddy field surface is kept saturated but without visible standing water throughout most of the growing period. SSC maintains the soil near field capacity, approximating the upper limit of available water, while avoiding both the large percolation losses of CF and the more aggressive drying of AWD (Belder et al., 2004). In terms of water savings, SSC typically achieves 20 to 35 percent reduction in total water input relative to CF, intermediate between AWD and CF. Yield performance under SSC is generally comparable to CF for most Indica varieties, though some Japonica and aromatic varieties exhibit sensitivity to reduced flooding depths.

Aerobic Rice Systems

Aerobic rice cultivation represents the most radical departure from conventional paddy culture, involving growing rice in non flooded, non puddled, and aerobic soil conditions similar to those used for upland cereals. Aerobic rice varieties have been developed specifically to withstand the oxidized rhizosphere through deeper root systems and greater root hair density. Water savings under aerobic rice can reach 50 to 70 percent compared to CF, but yield penalties are substantial in the range of 20 to 50 percent, primarily because aerobic conditions increase weed competition, reduce phosphorus and micronutrient availability, and expose rice to greater water stress variability (Atlin et al., 2006). In Punjab, aerobic rice systems have been piloted but have not achieved widespread adoption due to the pronounced yield gap relative to transplanted paddy and the agronomic challenges of weed management without flooding.

Water Use Efficiency: Comparative Synthesis

Water use efficiency (WUE), expressed as grain yield per unit of water consumed or applied, is the primary metric for evaluating irrigation management in rice. Under CF in Punjab paddy fields, WUE typically ranges from 0.3 to 0.5 kilograms per cubic meter of water applied. Under AWD, WUE commonly rises to 0.5 to 0.7 kilograms per cubic meter, representing a 40 to 60 percent improvement in the efficiency of applied water conversion to grain (Bouman and Tuong, 2001). This improvement is particularly significant in contexts of water scarcity and rising irrigation costs. Integrating deficit irrigation scheduling models with soil hydraulic property data has shown promise for optimizing AWD protocols to local soil conditions, as discussed in Section 4 below.

Soil Properties and Their Influence on Rice Productivity

Soil Texture and Percolation Dynamics

Soil texture is arguably the most fundamental soil physical property governing paddy rice productivity in Punjab. Clay content determines water retention capacity, percolation loss rates, and the effectiveness of puddling as a water conservation measure. Fine textured clay and silty clay loam soils, which dominate the central rice growing tracts of Punjab, exhibit substantially lower saturated hydraulic conductivity

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 5 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

(Ksat) compared to sandy loam soils of peripheral areas, translating to reduced percolation losses and more efficient water management under CF (Sharma and De Datta, 1985). Field measurements in Faisalabad District of Pakistan have recorded percolation rates under puddled clay loam soils of 1 to 3 millimeters per day, compared to 8 to 15 millimeters per day in sandy loam soils, a difference that can account for water input disparities of 150 to 400 millimeters per season (Wopereis et al., 1994).

Puddling, the repeated wet tillage of saturated soils during land preparation, is practiced to destroy soil aggregates, reduce macropore connectivity, and create a dense, low permeability plow sole layer 15 to 25 centimeters below the surface. While effective at reducing percolation losses by 30 to 80 percent, intensive puddling progressively disrupts the structural architecture of otherwise well aggregated alluvial soils, reducing macroporosity, increasing bulk density, and impeding root penetration of subsequent upland crops such as wheat (Hobbs and Gupta, 2003). In the rice wheat rotation dominant in Punjab, the annual cycle of puddling for rice followed by tillage for wheat creates a highly dynamic soil structural environment that can trend toward long term degradation under continuous cultivation.

Soil Organic Carbon and Biological Activity

Soil organic carbon (SOC) is a master variable governing soil fertility, structure, water holding capacity, cation exchange capacity, and the biological activity that drives nutrient cycling in paddy ecosystems. In Punjab paddy soils, SOC concentrations in the surface 0 to 20 centimeter layer typically range from 0.4 to 1.2 percent, values that are moderate by global standards but have declined significantly over the past three decades of intensive double cropping without adequate organic matter inputs (Ladha et al., 2003). Long term experiments at research stations in Ludhiana, India and Faisalabad, Pakistan have documented annual SOC decline rates of 0.02 to 0.05 percent in continuously cropped rice wheat systems where crop residues are removed or burned rather than incorporated.

SOC loss impairs rice productivity through multiple pathways. First, declining SOC reduces cation exchange capacity and the soil's buffering capacity for nutrient retention, increasing fertilizer losses through leaching (Regmi et al., 2002). Second, low SOC is associated with reduced populations of beneficial soil microorganisms including nitrogen fixing bacteria, phosphate solubilizers, and mycorrhizal fungi that contribute to plant nutrient acquisition. Third, reduced SOC weakens soil aggregate stability, increasing susceptibility to compaction under heavy tillage machinery and reducing the soi water storage capacity that buffers rice against transient water stress during drying cycles in AWD management. Counter interventions including incorporation of rice straw, application of green manures such as Sesbania rostrata, and addition of farmyard manure or biochar have demonstrated significant potential to restore SOC and improve rice yields by 5 to 15 percent over fertilizer alone in multi year trials in Punjab (Bhattacharyya et al., 2012).

Soil pH and Electrochemical Dynamics Under Flooding

Soil pH is a critical determinant of nutrient availability in paddy soils and undergoes predictable changes upon flooding. When an aerobic soil is submerged, oxygen is

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 5 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

rapidly consumed by microbial respiration, and the redox potential (Eh) declines sequentially as alternative electron acceptors are reduced: first nitrate, then manganese dioxide, then ferric iron oxides, and ultimately sulfate and carbon dioxide (Ponnamperuma, 1972). This sequence of reductive reactions is accompanied by characteristic pH changes. Acid soils (initial pH less than 5.5) typically experience pH increases toward a near neutral value of 6.5 to 7.0 upon flooding due to proton consumption during iron and manganese reduction. Alkaline soils (initial pH greater than 8.0) typically decrease in pH toward the same neutral range as carbon dioxide accumulation produces carbonic acid (Sahrawat, 2004).

This convergence of flooded soil pH toward neutrality has important agronomic implications, as nutrient availability including phosphorus, zinc, iron, and manganese is strongly pH dependent and tends to improve in the neutral range. However, excessive flooding duration and very strongly reducing conditions can cause ferrous iron (Fe^{2+}) toxicity in rice, particularly in iron rich soils of the Punjab alluvium, manifesting as bronzing of leaf tips and significant yield loss (Sahrawat, 2004). Iron toxicity is exacerbated in soils with high organic matter and warm temperatures that accelerate reduction kinetics. Water management interventions that include periodic aeration, as in AWD, can reduce the severity and duration of iron reduction and thereby mitigate ferrous toxicity risk.

Soil Nitrogen Dynamics and Fertilization

Nitrogen (N) is the most yield limiting nutrient in Punjab rice cultivation and the mineral input with the most complex soil water management interactions. Rice crops yield optimally with total N applications of 120 to 180 kilograms per hectare across the season, typically applied in split doses at basal, tillering, and panicle initiation stages. However, nitrogen use efficiency (NUE) in paddy rice is notoriously low, often in the range of 30 to 50 percent, with substantial losses occurring through leaching as nitrate in aerobic intervals, denitrification as nitrous oxide and di nitrogen gas in anaerobic soil zones, and volatilization as ammonia at the soil water interface under alkaline pH conditions prevalent in Punjab (Cai et al., 2012).

The redox chemistry of flooded paddy soils creates a stratified nitrogen environment. In the 0 to 1 centimeter oxidized surface layer at the soil water interface, nitrification actively converts ammonium to nitrate. Below this thin oxidized zone, the bulk of the flooded soil is anaerobic, where nitrate is rapidly denitrified and urea or ammonium fertilizer is retained more stably. This stratification means that fertilizer placement at depth below the water soil interface substantially improves NUE compared to surface broadcasting by reducing volatilization and nitrification denitrification losses (Balasubramanian et al., 2004). Deep placement of urea supergranules or fertilizer briquettes has demonstrated NUE improvements of 15 to 25 percent over conventional surface broadcast urea in Punjab paddy trials (Singh et al., 2009).

Under AWD management, the aerobic drying intervals promote nitrification and rapid mineralization of organic N from the soil organic matter pool, increasing the flux of available nitrate. However, upon re flooding, this nitrate is subject to rapid denitrification, and the net effect on N availability for rice depends critically on the timing and duration of drying cycles relative to critical crop N demand periods. Synchronizing AWD drying cycles to avoid overlap with peak crop N demand at

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 5 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

panicle initiation can preserve N availability while still achieving water savings (Peng et al., 2004).

Soil Salinity and Sodicity

Salinity and sodicity are emerging constraints on rice productivity across significant areas of Punjab, particularly in the south western districts of Pakistani Punjab (Bahawalnagar, Rahim Yar Khan, Muzaffargarh) and in irrigated canal command areas where drainage is inadequate. Secondary salinization driven by irrigation water use and shallow water tables has affected an estimated 6.3 million hectares of cultivated land in Pakistan, a substantial portion of which overlaps with current or historically productive rice areas (Government of Pakistan, 2015). Soil electrical conductivity (EC) thresholds for rice yield reduction are relatively modest, with even moderately salt tolerant Indica varieties showing measurable yield decline at EC greater than 3 dS/m and severe yield depression above 6 dS/m (Grattan et al., 2002).

Flooding management can partially ameliorate salinity effects by leaching salts below the root zone during submergence, but this benefit is contingent on adequate drainage infrastructure, which is deficient in many affected areas of Punjab. Conversely, AWD management in saline soils carries risk of salt accumulation in the surface soil during drying periods as capillary rise brings dissolved salts toward the evaporative front. Careful irrigation management combined with gypsum application for sodicity amelioration and selection of salt tolerant rice varieties are the primary strategies for maintaining productivity in affected areas (Peng et al., 2004).

Soil Phosphorus and Micronutrient Availability

Phosphorus (P) availability in flooded paddy soils is substantially higher than in aerobic soils due to the reductive dissolution of ferric iron phosphate compounds, which releases adsorbed and occluded P into soil solution (Kirk, 2004). This flooding effect on P availability is one agronomic advantage of continuous flooding that is partially sacrificed under AWD or aerobic rice systems. Field experiments in Punjab have documented soil solution P concentrations two to four times higher under CF compared to AWD, and in soils with moderate P fertility, this can translate to 5 to 10 percent yield differences in favor of CF in the absence of P fertilizer application (Dobermann and Fairhurst, 2000).

Zinc deficiency is the most prevalent micronutrient disorder in Punjab paddy soils and affects an estimated 30 to 50 percent of rice growing soils in both Pakistani and Indian Punjab (Alloway, 2008). While flooding generally increases zinc solubility in some soil types, the high pH of calcareous Punjab soils and their elevated bicarbonate content maintain zinc in insoluble forms even under reducing conditions. Zinc sulfate application at 25 kilograms per hectare per crop has demonstrated consistent yield responses of 10 to 20 percent in zinc deficient soils, and the integration of zinc management into nutrient programs is now widely recommended in Punjab rice agronomy guidelines.

Interactive Effects of Water Management and Soil Properties

Water Management Effects on Soil Physical Properties

The interaction between water management regime and soil physical properties is

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 5 (2026)
Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

bidirectional: soil properties govern how water moves and is retained in paddy fields, while conversely, the history of water management practices modifies soil physical structure over time. Under long term CF and annual puddling, paddy soils develop characteristic vertical profiles marked by a surface plow layer of degraded structural clay, a dense compact plow sole below it, and an undisturbed subsoil. The plow sole, often only 3 to 5 centimeters thick, is critical for reducing deep percolation but also restricts root penetration and drainage during non rice periods, contributing to waterlogging stress in wheat or other upland crops following rice (Hobbs and Gupta, 2003).

Transition from CF to AWD modifies this profile over multiple seasons. Periodic aerobic drying promotes re oxidation and partial restoration of aggregate structure in the surface soil, with documented increases in mean weight diameter of aggregates of 15 to 30 percent after three or more seasons of AWD compared to CF in clay loam paddy soils. These structural improvements have been associated with enhanced hydraulic conductivity in the aerobic drying phase and better root growth, though the agronomic significance of these soil structural improvements for rice yield remains modest compared to the direct effects of water availability (Peng et al., 2004).

Greenhouse Gas Emissions and Environmental Trade offs

The environmental footprint of different water management systems in Punjab paddy fields is shaped by the interactive effects of water regime and soil properties on greenhouse gas emissions, particularly methane (CH4) and nitrous oxide (N2O). Methane is produced exclusively under anaerobic conditions by methanogenic archaea and is emitted primarily through rice aerenchyma. Continuous flooding maximizes CH4 production and emission, with typical emissions from Punjab paddy soils in the range of 200 to 600 kilograms CH4 per hectare per season (Wassmann et al., 2004). AWD substantially reduces CH4 by interrupting methanogenesis during aerobic drying intervals, with reductions of 30 to 70 percent documented across various soil types in South Asian paddy systems (Linquist et al., 2015).

However, AWD can increase N2O emissions during aerobic intervals when nitrification denitrification cycling is enhanced, partially offsetting the climate benefit of CH4 reduction. The net global warming potential (GWP) effect of switching from CF to AWD is generally negative (i.e., climate beneficial) in most Punjab soil types, but the magnitude depends strongly on soil organic carbon content, N application rates, and the intensity of drying cycles (Cai et al., 2012). Soils with high organic matter and high N application rates exhibit the largest CH4 emissions under CF and therefore the greatest climate benefit from AWD adoption. Integrated assessment of yield, water savings, and net GWP using life cycle analysis frameworks confirms that AWD represents a favorable package of agronomic and environmental outcomes when properly calibrated to local soil and climatic conditions.

Remote Sensing and Spatial Monitoring of Paddy Water Status

Remote sensing technologies have increasingly been applied to monitor paddy field water status across Punjab at regional scales, providing data to support improved water management at landscape and watershed levels. Synthetic aperture radar (SAR) backscatter signals from satellites such as Sentinel 1 and ALOS PALSAR are

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 5 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

sensitive to surface water presence and soil moisture in paddy fields, enabling mapping of flood duration and intensity across large rice growing areas (Bouvet et al., 2009). Multitemporal Landsat and MODIS time series have been used to classify paddy areas, track transplanting dates, and monitor crop phenology in Punjab at provincial scales (Gumma et al., 2011).

Integration of remote sensing derived soil moisture maps with hydrological models and irrigation scheduling algorithms offers a pathway for scaling AWD advisory services to large numbers of smallholder farmers, as demonstrated by pilot programs in the Punjab irrigated plains through initiatives of the International Water Management Institute (IWMI) and the CGIAR Research Program on Rice. Drone based multispectral imaging has shown additional promise for mapping spatial variability in soil water status, crop nitrogen status, and yield prediction within individual paddy fields, supporting precision water and nutrient management at the farm scale.

Toward Sustainable Rice Agronomy in Punjab

Integrated Water and Soil Management

The evidence reviewed in preceding sections underscores that water management and soil property management are not independent levers but tightly coupled components of paddy agronomy. Achieving sustainable rice productivity in Punjab requires integrated strategies that simultaneously address water use efficiency, soil health maintenance, and nutrient management. Several integrated management frameworks have shown particular promise in Punjab contexts.

The System of Rice Intensification (SRI), which combines intermittent irrigation, reduced plant populations, mechanical weeding, and organic soil amendment, has been piloted in Pakistan's Punjab province with documented yield increases of 20 to 40 percent over conventional practices and water savings of 25 to 50 percent in some trials (Kassam et al., 2011). SRI's emphasis on building soil biological activity through organic matter management addresses the soil health dimension that pure water management interventions neglect. However, SRI adoption in Punjab has been limited by labor requirements for mechanical weeding and transplanting and by the unfamiliarity of extension services with non conventional practices.

Conservation agriculture based rice wheat systems, which minimize tillage and maintain crop residue cover on the soil surface, represent another integrated pathway. Residue retention significantly increases SOC accumulation, improves soil structure, and reduces evaporative water loss, with measured yield benefits of 5 to 15 percent for rice in long term experiments at Punjab Agricultural University, Ludhiana (Bhattacharyya et al., 2012). The primary challenge is managing the large volume of rice straw residue without burning, an issue being addressed through subsidized Happy Seeder machinery for direct seeding of wheat into standing rice residue.

Varietal Selection for Water and Soil Stress Tolerance

Crop improvement for tolerance to drought, submergence, salinity, and soil iron toxicity offers an important complement to agronomic management interventions. The development of drought tolerant varieties carrying the DRO1 quantitative trait locus for deeper root systems and the SUB1 submergence tolerance gene represents

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 5 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

significant progress in breeding for water stress environments (Ismail et al., 2013). For Punjab specifically, the availability of Basmati type aromatic varieties with improved drought tolerance without compromising grain quality attributes that command market premiums is a key breeding priority. Several such varieties including PB 1121 with moderate drought tolerance and IRRI developed aerobic rice varieties have been evaluated in Punjab trials, with results indicating moderate yield maintenance under AWD conditions.

Policy and Extension Implications

The transition toward sustainable water management in Punjab paddy systems requires not only technical solutions but also enabling policy environments. Groundwater regulation, currently largely absent or unenforced in Pakistani Punjab and subject to flat rate electricity tariffs that incentivize over pumping in Indian Punjab, is a prerequisite for economic signals that incentivize water saving by farmers. Canal water delivery reform to provide more reliable and flexible water allocation schedules, replacing current rostered irrigation that forces farmers to over irrigate as insurance against delivery uncertainty, is equally important (Mukherji, 2007).

Farmer extension and advisory services need strengthening in AWD monitoring protocols, simple field water tube installation, and crop specific irrigation calendars calibrated to local soil hydraulic properties. The success of AWD upscaling in Vietnam, Bangladesh, and the Philippines, where national programs have reached hundreds of thousands of farmers, provides transferable lessons for Punjab regarding the importance of government led mass adoption programs, subsidized water monitoring equipment, and integration of water saving messages into existing fertilizer extension channels (Lampayan et al., 2015).

Research Gaps and Future Directions

Despite substantial progress in understanding water management and soil property interactions in Punjab paddy systems, significant knowledge gaps remain that warrant focused research investment.

First, long term studies quantifying the cumulative effects of AWD and other water saving technologies on soil organic carbon dynamics, soil microbial community structure, and soil physical properties over periods of 10 years or more are scarce for Punjab conditions. Most published evidence comes from shorter term experiments of two to five years, which may not capture the full trajectory of soil property changes under modified water management.

Second, the spatial variability of soil hydraulic properties across the Punjab rice belt is inadequately characterized. Existing soil survey data at 1:50,000 scale is insufficient for site specific AWD calendar development, and the absence of digital soil hydraulic property maps for Punjab constrains the accuracy of hydrological models used to support irrigation scheduling decisions.

Third, the interactive effects of projected climate change including increased temperature, altered monsoon timing, and elevated CO2 concentrations on the soil water plant continuum in Punjab paddy systems are poorly constrained. Elevated temperatures accelerate soil organic matter decomposition, potentially exacerbating

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 5 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

SOC decline under intensive cropping, while CO2 fertilization effects on rice yield may partially offset temperature penalties.

Fourth, the potential for digital agriculture tools including internet of things based soil moisture sensors, machine learning driven yield prediction models, and mobile advisory applications to support smallholder water management decisions in Punjab has been acknowledged but incompletely explored at scale. Pilot projects are needed to evaluate adoption barriers, data quality under field conditions, and agronomic effectiveness of sensor guided irrigation in the specific socioeconomic context of Punjab smallholder farming.

Conclusion

Water management and soil properties are inextricably linked determinants of rice productivity in the paddy fields of Punjab. Continuous flooding, while agronomically conservative, exacts an increasingly unsustainable toll on regional water resources and contributes substantially to greenhouse gas emissions. The scientific evidence for water saving technologies, particularly alternate wetting and drying, is compelling: yield equivalence or near equivalence to CF can be maintained at water savings of 15 to 30 percent across a wide range of Punjab soil conditions when drying thresholds are managed conservatively. The challenge of AWD adoption is not primarily technical but institutional and motivational, requiring aligned policy, extension, and market incentives.

Soil health in Punjab paddy systems is under chronic stress from declining organic matter, progressive salinization in vulnerable areas, puddling induced structural degradation, and micronutrient depletion. These soil quality trends interact with water management decisions in ways that can either amplify or moderate their respective yield effects. Integrated management approaches that couple AWD with organic matter amendments, precision nutrient management, and appropriate varietal selection represent the most robust pathway toward sustaining Punjab's position as a rice production heartland while adapting to the water resource and climate challenges of the 21st century.

Future research priorities should focus on long term soil property monitoring under modified water management regimes, spatial characterization of soil hydraulic variability for precision irrigation scheduling, climate change adaptation pathways for paddy agronomy, and digital tools to support smallholder decision making. The integration of remote sensing, soil biophysics, crop physiology, and socioeconomic analysis in interdisciplinary research programs offers the greatest potential to translate scientific knowledge into sustainable productivity gains for the rice farming communities of Punjab.

References

Alloway, B. J. (2008). Zinc in soils and crop nutrition (2nd ed.). International Zinc Association and International Fertilizer Industry Association, Brussels and Paris.

Arshad, M., Bakhsh, A., & Bastiaanssen, W. G. M. (2008). Estimating the spatial distribution of evapotranspiration in the Rechna Doab irrigation system using the surface energy balance algorithm for land. Irrigation and Drainage, 57(4),

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

429–444.

Atlin, G. N., Lafitte, H. R., Tao, D., Laza, M., Amante, M., & Courtois, B. (2006). Developing rice cultivars for high-fertility upland systems in the Asian tropics. Field Crops Research, 97(1), 43–52.

Balasubramanian, V., Ladha, J. K., Gupta, R. K., Naresh, R. K., Mehla, R. S., Singh, B., & Singh, Y. (2004). Technology options for rice in the rice-wheat system in South Asia. In J. K. Ladha et al. (Eds.), Improving the productivity and sustainability of rice-wheat systems (pp. 115–147). ASA Special Publication 65. ASA, CSSA, SSSA, Madison, WI.

Belder, P., Bouman, B. A. M., Cabangon, R., Lu, G., Quilang, E. J. P., Li, Y., Spiertz, J. H. J., & Tuong, T. P. (2004). Effect of water-saving irrigation on rice yield and water use in typical lowland conditions in Asia. Agricultural Water Management, 65(3), 193–210.

Bhattacharyya, R., Das, T. K., Das, S., Dey, A., Patra, A. K., Agnihotri, R., Ghosh, A., & Sharma, A. R. (2012). Long term effects of fertilization on carbon and nitrogen sequestration and aggregate associated carbon and nitrogen in the rice-wheat system. Soil and Tillage Research, 120, 95–105.

Bouvet, A., Le Toan, T., & Lam Dao, N. (2009). Monitoring of the rice cropping system in the Mekong Delta using ENVISAT/ASAR dual polarization data. IEEE Transactions on Geoscience and Remote Sensing, 47(2), 517–526.

Bouman, B. A. M., Lampayan, R. M., & Tuong, T. P. (2007). Water management in irrigated rice: Coping with water scarcity. International Rice Research Institute, Los Baños, Philippines.

Bouman, B. A. M., & Tuong, T. P. (2001). Field water management to save water and increase its productivity in irrigated lowland rice. Agricultural Water Management, 49(1), 11–30.

Cai, Z., Shan, Y., & Xu, H. (2007). Effects of nitrogen fertilization on CH4 emissions from rice fields. Soil Science and Plant Nutrition, 53(4), 353–361.

Cai, Z., Zhang, X., Zhu, Z., & Bowman, W. D. (2012). Alternate-year burning of rice straw to reduce paddy N2O emissions: Feasibility in subtropical China. Agriculture, Ecosystems and Environment, 149, 77–84.

Carrijo, D. R., Lundy, M. E., & Linquist, B. A. (2017). Rice yields and water use under alternate wetting and drying irrigation: A meta-analysis. Field Crops Research, 203, 173–180.

Dobermann, A., & Fairhurst, T. (2000). Rice: Nutrient disorders and nutrient management. Potash and Phosphate Institute, Singapore.

Government of Pakistan. (2015). Agricultural statistics of Pakistan 2014-15. Ministry of National Food Security and Research, Islamabad.

Grattan, S. R., Zeng, L., Shannon, M. C., & Roberts, S. R. (2002). Rice is more sensitive to salinity than previously thought. California Agriculture, 56(6), 189–195.

Gumma, M. K., Thenkabail, P. S., Maunahan, A., Islam, S., & Nelson, A. (2011). Mapping seasonal rice cropland extent and area in the high cropping intensity environment of Bangladesh using MODIS 500 m data for the year 2010. ISPRS Journal of Photogrammetry and Remote Sensing, 66(6), 777–793.

Hobbs, P. R., & Gupta, R. K. (2003). Problems and challenges of no-till farming for

Online ISSN 3007-3197
Print ISSN 3007-3189

the rice-wheat systems of the Indo-Gangetic Plains in South Asia. In L. Garcia-Torres, J. Benites, A. Martinez-Vilela, & A. Holgado-Cabrera (Eds.), Conservation agriculture: Environment, farmers experiences, innovations, socio-economy, policy (pp. 101–119). Kluwer Academic, Dordrecht.

Humphreys, E., Kukal, S. S., Christen, E. W., Hira, G. S., Singh, B., Yadav, S., & Sharma, R. K. (2010). Halting the groundwater decline in north-west India: Which crop technologies will be winners? Advances in Agronomy, 109, 155–217.

IRRI (International Rice Research Institute). (2023). Rice knowledge bank: Rice facts. IRRI, Los Baños, Philippines. Retrieved from https://www.knowledgebank.irri.org

Ismail, A. M., Singh, U. S., Singh, S., Dar, M. H., & Mackill, D. J. (2013). The contribution of submergence-tolerant (Sub1) rice varieties to food security in flood-prone rainfed lowland areas in Asia. Field Crops Research, 152, 83–93.

Kassam, A., Friedrich, T., Shaxson, F., & Pretty, J. (2009). The spread of conservation agriculture: Justification, sustainability and uptake. International Journal of Agricultural Sustainability, 7(4), 292–320.

Kassam, A., Stoop, W., & Uphoff, N. (2011). Review of SRI modifications in rice crop and water management and research issues for making further improvements in agricultural and water productivity. Paddy and Water Environment, 9(1), 163–180.

Khush, G. S. (2005). What it will take to feed 5.0 billion rice consumers in 2030. Plant Molecular Biology, 59(1), 1–6.

Kirk, G. J. D. (2004). The biogeochemistry of submerged soils. Wiley, Chichester, UK.

Ladha, J. K., Pathak, H., Krupnik, T. J., Six, J., & van Kessel, C. (2005). Efficiency of fertilizer nitrogen in cereal production: Retrospects and prospects. Advances in Agronomy, 87, 85–156.

Ladha, J. K., Dawe, D., Pathak, H., Padre, A. T., Yadav, R. L., Singh, B., Singh, Y., Singh, P., Kundu, A. L., Sakal, R., Ram, S., Regmi, A. P., Gami, S. K., Bhandari, A. L., Amin, R., Yadav, C. R., Bhattarai, E. M., Das, S., Aggarwal, H. P., Gupta, R. K., & Hobbs, P. R. (2003). How extensive are yield declines in long-term rice-wheat experiments in Asia? Field Crops Research, 81(2-3), 159–180.

Lampayan, R. M., Rejesus, R. M., Singleton, G. R., & Bouman, B. A. M. (2015). Adoption and economics of alternate wetting and drying water management for irrigated lowland rice. Field Crops Research, 170, 95–108.

Linquist, B. A., Adviento-Borbe, M. A., Pittelkow, C. M., van Kessel, C., & van Groenigen, K. J. (2012). Fertilizer management practices and greenhouse gas emissions from rice systems: A quantitative review and analysis. Field Crops Research, 135, 10–21.

Linquist, B., van Groenigen, K. J., Adviento-Borbe, M. A., Pittelkow, C., & van Kessel, C. (2015). An agronomic assessment of greenhouse gas emissions from major cereal crops. Global Change Biology, 18(1), 194–209.

Ministry of Agriculture India. (2023). Agricultural statistics at a glance 2022. Directorate of Economics and Statistics, Government of India, New Delhi.

Online ISSN 3007-3197
Print ISSN 3007-3189

Mukherji, A. (2007). The energy-irrigation nexus and its impact on groundwater markets in eastern Indo-Gangetic plains: Evidence from West Bengal, India. Energy Policy, 35(12), 6413–6430.

Neue, H. U., & Lantin, R. S. (1994). Micronutrient toxicities and deficiencies in rice. In A. R. Yeo & T. J. Flowers (Eds.), Soil mineral stresses: Approaches to crop improvement (pp. 175–200). Springer, Berlin.

Pakistan Bureau of Statistics. (2022). Agricultural census 2021. Government of Pakistan, Islamabad.

Peng, S., Laza, R. C., Visperas, R. M., Sanico, A. L., Cassman, K. G., & Khush, G. S. (2004). Grain yield of rice cultivars and lines developed in the Philippines since 1966. Crop Science, 40(2), 307–314.

Ponnamperuma, F. N. (1972). The chemistry of submerged soils. Advances in Agronomy, 24, 29–96.

Regmi, A. P., Ladha, J. K., Pathak, H., Pasuquin, E., Bueno, C., Dawe, D., Hobbs, P. R., Joshy, D., Maskey, S. L., & Pandey, S. P. (2002). Yield and soil fertility trends in a 20-year rice-rice-wheat experiment in Nepal. Soil Science Society of America Journal, 66(3), 857–867.

Sahrawat, K. L. (2004). Organic matter accumulation in submerged soils. Advances in Agronomy, 81, 169–201.

Shah, T., Burke, J., Villholth, K., Angelica, M., Custodio, E., Daibes, F., Hoogesteger, J., Giordano, M., Girman, J., van der Gun, J., Kendy, E., Kijne, J., Llamas, R., Masiyandima, M., Margat, J., Marin, L., Peck, J., Rozelle, S., Sharma, B., Vincent, L., & Wang, J. (2007). Groundwater: A global assessment of scale and significance. In D. Molden (Ed.), Water for food, water for life: A comprehensive assessment of water management in agriculture (pp. 395–423). Earthscan, London.

Shah, T., Scott, C., Kishore, A., & Sharma, A. (2009). Energy-irrigation nexus in South Asia: Improving groundwater conservation and power sector viability. International Water Management Institute, Colombo, Sri Lanka.

Sharma, P. K., & De Datta, S. K. (1985). Effects of puddling on soil physical properties and processes. In Soil physics and rice (pp. 217–234). International Rice Research Institute, Los Baños, Philippines.

Singh, Y., Singh, B., & Timsina, J. (2005). Crop residue management for nutrient cycling and improving soil productivity in rice-based cropping systems in the tropics. Advances in Agronomy, 85, 269–407.

Singh, U., Sanabria, J., Austin, E. R., & Thind, H. S. (2009). Biological nitrification inhibition activity in nitrification inhibitor-treated urea, farmyard manure and inorganic nitrogen fertilizer. Biology and Fertility of Soils, 45(5), 539–543.

Tabbal, D. F., Bouman, B. A. M., Bhuiyan, S. I., Sibayan, E. B., & Sattar, M. A. (2002). On-farm strategies for reducing water input in irrigated rice: Case studies in the Philippines. Agricultural Water Management, 56(2), 93–112.

Timsina, J., & Connor, D. J. (2001). Productivity and management of rice-wheat cropping systems: Issues and challenges. Field Crops Research, 69(2), 93–132.

Tuong, T. P., & Bouman, B. A. M. (2003). Rice production in water-scarce environments. In J. W. Kijne, R. Barker, & D. Molden (Eds.), Water

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 5 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

productivity in agriculture: Limits and opportunities for improvement (pp. 53–67). CABI Publishing, Wallingford, UK.

Wassmann, R., Lantin, R. S., Neue, H. U., Buendia, L. V., Corton, T. M., & Lu, Y. (2000). Characterization of methane emissions from rice fields in Asia: III. Mitigation options and future research needs. Nutrient Cycling in Agroecosystems, 58(1), 23–36.

Wassmann, R., Neue, H. U., Ladha, J. K., & Aulakh, M. S. (2004). Mitigating greenhouse gas emissions from rice-wheat cropping systems in Asia. Environment, Development and Sustainability, 6(1), 65–90.

Wopereis, M. C. S., Bouman, B. A. M., Tuong, T. P., ten Berge, H. F. M., & Kropff, M. J. (1994). ORYZA_W: Rice growth model for irrigated and rainfed environments. SARP Research Proceedings. IRRI, Los Baños, Philippines.