Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/a

Climate Change-Driven Microbial Dynamics in Food Systems: Implications for Pathogen Emergence, Food Safety, and Global Human Health

Dr. Abdullah*, Fouzia Hussain**, Razia Kalsoom**, Dr. Tahseen Aslam**, Tanveer Ahmad Khan**, Waseem Aslam, Muhammad Afzal**, Dn. Sheema Nauman***
* Corresponding Author, PCSIR Labs, Islamabad. Email: abdullahese@gmail.com,
**Co-Author, PCSIR Labs, Islamabad
***Co-Author, Aaymed Care Medical Centre, Minsa Hospital, Rawalpindi

Paper Received (25th January 2026), 1st Review (30th March 2026), 2nd Review (15th April 2026), Paper Accepted (30th April 2026), Paper Published (16th June 2026)

ABSTRACT

Climate change is fundamentally reshaping microbial communities throughout the global food supply chain, driving the emergence and re-emergence of foodborne pathogens with profound consequences for food safety and human health. Rising temperatures, altered precipitation patterns, extreme weather events, and sea-level rise accelerate microbial metabolism, stress adaptation, biofilm formation, and horizontal gene transfer, expanding the geographic and seasonal range of key bacteria such as Salmonella, Campylobacter, Vibrio, and Shiga toxin-producing E. coli, as well as norovirus, protozoan parasites, and mycotoxin-producing fungi. These climate-driven dynamics intensify contamination risks at every stage from drought-stressed irrigation and flood-mobilized runoff in production fields to disrupted cold chains in processing, storage, and distribution while promoting zoonotic spillover, antimicrobial resistance, and supply-chain vulnerabilities. The

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)
Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/a

result is a sharply rising global disease burden that disproportionately affects children, the elderly, immune-compromised individuals, and low-income regions, with stark regional disparities across Africa, Asia, Europe, and the Americas and trillions in combined economic and public-health costs. Observational studies, modeling projections, and field evidence confirm that specific climate variables directly correlate with increased incidence and outbreak severity. Practical mitigation and adaptation strategies including climate-informed HACCP and Quantitative Microbial Risk Assessment, climate-smart agriculture, enhanced One-Health surveillance with whole-genome sequencing and predictive modeling, targeted education, and strengthened international policy cooperation offer scalable solutions. Addressing remaining gaps in longitudinal data, mechanistic research, regional models, and equity considerations is essential. Urgent, coordinated global action is required to safeguard food safety and public health in a rapidly warming world.

Keywords: **Climate change, foodborne pathogens, microbial dynamics, pathogen emergence, food safety, One Health**

Introduction

Global food systems are undergoing fundamental transformations due to the climate change by increasing warming, changing the patterns of precipitation, and the frequency of extreme weather events. The Sixth Assessment Report Synthesis by the Intergovernmental Panel on Climate Change has shown that the global surfaces have already experienced an increase of about 1.1 o C above the pre-industrial temperature, with estimates indicating high chances of exceeding 1.5 o C by 2021 to 2040 under the current trends (IPCC, 2023). Such changes are in the form of increased intensity of heatwaves, long-lasting droughts in the subtropical world, increased occurrence of heavy rainfalls, and rise of the sea level, which directly affect agricultural productivity, water supply, and stability of the ecosystem. Such drivers in food systems enhance farm-to-fork vulnerabilities: high temperatures enhance microbial metabolism and stress responses, and changed precipitation patterns augment flooding and drought cycles that mobilize contaminants in soil, water, and livestock habitats. These impacts multiply with each increase in 0.5C of warming,

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)
Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/a

resulting in noticeable increases in heat extremes, heavy precipitation, and regional droughts that impact crop yields and livestock health in low- and mid-latitude areas, where food insecurity is already acute (World Resources Institute, 2023). More importantly, these climatic changes do not exist in vacuums rather they combine with microbial communities to form new ecological niches, which are preferent to growth of opportunistic pathogens and a change in community structure that the traditional food safety paradigm has yet to fully account.

The microbial dynamics of food systems is a complicated interaction of development, existence, adjustment, group changes, and horizontal gene exchange all of which are significantly affected by these climate drivers. The soil, water, crops, livestock, and processing environment microorganisms are sensitive to environmental changes: an increase in temperature stimulates metabolic activity and biofilm development, allowing pathogens like Salmonella, Campylobacter, and Vibrio species to multiply and spread more effectively, whereas changes in humidity and extreme events stimulate horizontal gene transfer of virulence and antibiotic-resistance factors (Naga, 2025). The alteration of community at the community level is through the selection of resilient taxa by climate stressors that replaces beneficial microbes and establishes dysbiotic conditions, which increase the probability of contamination during harvest, storage, and distribution. As an example, compaction of the soil due to drought, and flooding occurrences promote the movement of enteric bacteria in animal waste to irrigation sources and produce fields, and ocean warming introduces new coastal harvesting areas to Vibrio vulnificus (Toor, 2024). These processes are not just ecological oddities but have a direct role in the rise and fall of foodborne pathogens by increasing geographic coverage, broadening seasonal periods of transmission and increasing virulence due to genetic adaptations occurring under stress. Analytically, this is a self-perpetuating feedback cycle whereby the fact that climate exacerbates microbial resilience is taking down the pillars of food safety, turning formerly stable supply chains into piping systems of contamination.

These microbial changes have an epidemiological footprint that is grim and measurable, and is connected to a continuing global foodborne disease burden that is

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)
Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/a

attributable to climate changes. According to the estimates of the World Health Organization, unsafe food makes 1 out of 10 individuals in the world get ill each year, which is estimated as 600 million cases, 420,000 deaths, and 33 million healthy life years, with the number of children under the age of five contributing almost a third of the fatalities (World Health Organization, n.d.). Major pathogens other than Toxoplasma cause an estimated 9.9 million domestically acquired illnesses, 53,300 hospitalizations and 931 deaths annually in the United States alone with produce, meat, poultry, and dairy being the main vehicles (Centers for Disease Control and Prevention, 2025). This trend is supported by regional data: the increase in temperatures is associated with an increase in the incidence of Salmonella and Campylobacter in subtropical and temperate regions and extreme events have caused reported outbreaks by contaminated water and flooded fields (Awad et al., 2024). This is a review that, systematically, is looking at these mechanisms driven by climate across the complete farm-to-fork continuum, how microbial adaptation and community restructuring leads to increased pathogen emergence, compromised food safety and adverse human-health effects. Search of literature: MEDLINE (PubMed), Web of Science, Scopus and Google Scholar were searched thoroughly to identify articles and reports in English that included keywords such as climate change, foodborne pathogens, microbial dynamics, food chain, farm to fork, emergence and food safety and which were published between 2000 and 2024. Relevance based screening was performed on titles, abstracts and full texts and preferred empirical studies, systematic reviews and authoritative syntheses of climate-microbe-food-health linkages which offer mechanistic or quantitative evidence. The synthesis of these understandings highlights in the review that there is an urgent demand to implement integrated and climate-resilient interventions to protect the world food systems and human health.

The Link between Climate Change and Microbial Dynamics in Food Systems

An increase in global temperature and changes in humidity are effective drivers of microbial metabolism and stress reactions in food production systems, which fundamentally changes the fitness of microbiomes living in soil to food storage after harvest. The warmer temperatures increase the speed of enzymes, replication in

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)
Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/a

bacteria like Salmonella and Escherichia coli, which leads to a rapid increase in population in crops and animals and a spike in humidity results in micro environments that extend the lifespan on plant surfaces and in irrigation water. Such changes initiate adaptive stress, such as increasing heat-shock proteins which improve thermal tolerance and osmotic regulation systems which oppose desiccation during drought pulses. Such dynamics, analytically, provide a selective pressure in which even small changes in temperature of 1-2 C can reduce lag phases, and extend exponential growth and turn stable commensal communities into opportunistic contaminants that endanger food safety at numerous chain stages.

Severe weather conditions and rising sea levels add enormous bursts of contamination to food systems overpowering the natural defenses and spreading pathogens by floodwaters, stormwater runoff, and infrastructure collapse. The process of hurricanes, heavy rains, and inundation of the coast places the fecal matter of livestock activities and sewerage systems into fields of farming and processing facilities and provides the conditions of high nutrient and stagnation that supports the growth of explosive bacteria. Not only do these events carry along pathogens, but also cause physiological stress, which increases the virulence gene expression of surviving populations, and salinizes freshwater aquifers and estuaries due to sea-level intrusion and thereby selects halotolerant species that can survive longer in brackish irrigation sources. The impact is a compounding risk where disrupted cold chains in periods of power failure caused by storms increase spoilage, and recovery operations after the event can frequently be dependent on spoiled water supplies, introducing contamination deep into the supply chains and increasing the outbreak potential many times beyond the storm affected disaster area.

Climate-mediated range expansion of foodborne pathogens also causes new areas and crops to be exposed to formerly exotic microbes, and rapid selection of adaption in quorum sensing, biofilm formation, and expression of virulence genes. With the expansion of favorable conditions northward and up the mountain, the pathogen such as Campylobacter and emerging strains inhabit new farming areas, where precipitation conditions are shifting, allowing horizontal gene transfer of the resistance and virulence factors. On the community level, the quorum sensing

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)
Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/a

organizes community-level behavior, allowing the production of biofilm matrix on equipment surfaces and produce that protects populations against sanitizers and other environmental stressors. This ecological niche engineering This method of adaptive restructuring analytically increases persistence during transport and storage, and modeling forecasts that up to 200 percent of current incidences will increase by century end under moderate warming conditions in northern latitudes.

The ecological impact of ocean and coastal warming radically rearranges marine microbial communities, widening both the geographic and seasonal range of Vibrio species in food chains, and increasing food safety risks by increasing biofilm resistance and virulence. High sea-surface temperatures (more than 20C) stimulate the rapid growth of Vibrio vulnificus and V. parahaemolyticus in shellfish and finfish and extreme events, such as heatwaves and storms, further spread these opportunists to brackish harvesting areas. Density-dependent signaling then regulates biofilm formation on shellfish surfaces and processing equipment, which give the shellfish resistance to depuration and cold storage, as well as support viable but non-culturable states that reactivate on consumption. This is a climate-enhanced feedback mechanism: as waters grow warmer and stratified, the amount of oxygen and the proportion of pathogenic genotypes increase, forecasting dramatic increases in cases of vibriosis in Europe and North America, as globalization of seafood trade spreads these hazards throughout the world and weakens traditional control methods based on the stability of ocean conditions.

3. Mechanisms Driving Pathogen Emergence and Re-Emergence

Climate change exerts severe thermal and osmotic stresses that induce microbial evolution and resistance to stress provide genetic and physiological capabilities allowing foodborne pathogens to survive in derailed farm-to-fork conditions. In bacteria like Salmonella and Escherichia coli, in response to increasing temperatures, heat-shock proteins and efflux pumps that release toxins and preserve membrane integrity are activated, and long drought cycles are also a selective pressure favoring those bacteria that can enter viable but non-culturable states to be reactivated on rehydration. Those mutations increase the transfer of resistance cassettes horizontally, transforming populations that were initially vulnerable into hardy

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)
Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/a

survivors that can live longer on produce surfaces and in irrigation systems. The outcome is the increase of contamination incidences as microorganisms take advantage of climatic stressed niches, where high metabolic rates decrease the generation time and increase toxin production despite refrigerated supply chains. This resilience is the direct cause of the re-emergence of pathogens that are now being checked by food safety measures with relative ease.

Changing climatic patterns enhance zoonotic spillover and host-range expansion through forcing wildlife, livestock and humans into previously unseen ecological coexistence in food production environments. Shifting rain and warmer winters increase the length of the activity seasons of hosts like rodents and birds living in reservoirs, increasing the ability of such pathogens like Campylobacter and new strains of viruses to cross species barriers at the livestock-wildlife interface. These spillovers increase the host ranges because vectors move polewards and to new agricultural areas, bringing with them microbes that rapidly adapt to new mammalian or avian immune systems due to the high rate of mutation accelerated by environmental stress. This dynamic in poultry and cattle production generates new reservoirs that are contaminated with meat and dairy in the course of slaughter and milking, entrenching new strains in the world supply chains. The process converts localized enzootic cycles to endemic foodborne hazards that affect the current surveillance systems.

Increased food production under the stress of climate also increases the rate of pathogen outbreak due to increased management shortcuts that promote the growth of microbes throughout the chain of supply. The heatwaves and unpredictable rainfall make farmers to increase their use of antibiotics in livestock and use higher stocking rates, which select multidrug-resistant strains and allow their introduction into processing facilities. Droughts involving irrigation with recycled or brackish water increases the concentration of contaminants, and the decontamination of food crops is stressful with shorter harvest-to-market periods, thus decreasing the points of critical control. The changes in operations intensify the adaptation of microbes and Zoonotic threats and cause self-perpetuating contamination cycles.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/a



As illustrated in Figure 1, a conceptual model maps climate drivers (rising temperatures, extreme precipitation, sea-level rise) to microbial shifts (adaptation, community restructuring, virulence upregulation) and onward to pathogen emergence, with arrow thickness scaled to the strength of supporting evidence from field and modeling studies. The framework reveals how these mechanisms converge to erode food safety margins and elevate human exposure worldwide.

4. Impact on Specific Foodborne Pathogens

4.1 Foodborne Bacteria

Salmonella spp. exhibit pronounced relationships with increasing temperatures, and observations in Europe and Australia indicate that a 1 o C increase in weekly maximum temperature correlates with an 8.8 percent increase in cases, as a result of faster growth of pathogenic bacteria in contaminated foods in warmer periods. The highest seasonal increases are in summer where heat stimulates poultry, eggs and produce growth, and heavy rainfall and floods contribute to contamination of runoff of animal waste into irrigation sources and processing plants. Projection models show a geographic shift northward to areas that were cooler such as Scandinavia which may experience extended warm intervals which may increase incidence by mid-century due to intensified precipitation patterns. These climatic factors raise ambient temperatures coupled with humidity bursts in reduction of lag phase and augmented survival on surfaces, and this directly boosts outbreak prevalence in subtropical and temperate regions.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)
Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/a

The same applies to Campylobacter infections where ambient temperature and the previous heavy rainfall are powerful forecasts of human cases associated with poultry and poisoned water. In Germany and northern Europe, summer peaks prevail and prevalence is increased in broiler flocks and in indirect behaviors like barbecue which increases exposure. Extreme precipitation occurs, leading to outbreaks through the mobilization of fecal material in the surface waters, and drought decreases cases only to recover in the future through rehydration. Geographic changes are marked, and the models predict significant increases in the latitude of the north as the climate blankets and transforms what was formerly seasonal dangers into risks throughout the year in the broadened agricultural areas.

E. coli (STEC) associated with Shiga toxin, Vibrio spp., Listeria monocytogenes, Clostridium perfringens, Staphylococcus aureus, and Arcobacter have a similar climate sensitivity. The Vibrio species survive at temperatures above 1215C in coastal and brackish waters, and as the ocean warms and sea levels rise, they are extending their habitat to new shellfish harvesting regions in Europe and Alaska, where heat waves have already instigated documented expansion of seafood related diseases. Listeria takes advantage of temperature abuse of perishable foods such as ready-to-eat meat and dairy, and cold chains that are disrupted with climate changes during storms enhance persistence and growth even at refrigeration levels. The production of toxins by Clostridium perfringens and S. aureus is higher in warmer storage conditions and Arcobacter is more common in livestock carcasses in case of floods that undermine cleanliness. In these bacteria, recent observational studies in 2023-2025 associate specific variables temperature increases, heavy precipitation, and moisture with increased occurrence and modeling confirms poleward and altitude changes that have been problematic in both developed and developing areas to the current control strategies.

4.2 Foodborne Viruses

The spread of norovirus and hepatitis A virus aggravates whenever there is climate-induced flooding and change in temperatures which interfere with the water and sanitation systems. Wastewater systems are overwhelmed by flood occurrences, introducing viral particles in drinking water, fresh produce, and seafood supply, and

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)
Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/a

norovirus survives well in standing water and demonstrates an increased outbreak in the case of extreme precipitation. Seasonal trends show that temperatures are higher in the winter months in the temperate areas, but the modeling demonstrates that warmer and wetter world conditions may increase transmission periods throughout the year as the environment becomes more persistent and human crowding becomes more common in displacement situations. Geographic changes can be observed in those areas that have been experiencing increasing storms whereby post-harvest and irrigation processes expose viruses to global supply chains that amplify risks both in urban and rural environments.

Hepatitis A is transmitted in parallel with heavy rainfall and the rise in sea levels contaminating shellfish and produce by way of sewerage overflows, and thus the disease is more common in regions where sanitation infrastructure is insufficient. Observational studies also connect such events with the surges in waterborne transmission, and temperature increases during warmer seasons can stabilize viral particles on food surfaces. The interaction between the changed precipitation and extreme weather impose novel susceptibility in the production of the berry and leafy green where viruses survive longer in the changing humidity which directly correlates the climate variables with increased outbreak geography that goes beyond the traditional endemic regions.

4.3 Foodborne Parasites

Protozoan parasites like the Cryptosporidium, Cyclospora cayetanensis, Toxoplasma gondii and Giardia duodenalis take advantage of the change in water quality and host ecology brought about by climate to increase their foodborne footprint. The heavy rain and flooding moves oocysts of animal waste into surface water and irrigation systems with Cryptosporidium and Giardia demonstrating increased survival and transportation during extreme rain events which overflow into treatment facilities. These are appropriate seasonal trends in the temperate and subtropical areas are wetter and modeling predicts larger geographic distribution with warmer conditions favoring oocyst viability and pole-shifting in the host-vector interactions. The C. Difficile outbreaks in food have been associated with contaminated water sources during storms, and the Toxoplasma risk has increased

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)
Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/a

in the livestock exposed to the altered grazing habits in drought and heat stress, which entrench the parasites in deeper in the meat and fresh produce supply chains across the globe.

4.4 Mycotoxin-Producing Fungi and Other Emerging Microbes

Fungi producing mycotoxins, especially Aspergillus flavus (aflatoxins) and Fusarium species (deoxynivalenol, fumonisins), are very sensitive to higher temperatures and to changes in humidity and climate change is expected to cause more contamination in maize, peanuts and tree nuts in tropical and subtropical regions. The future perspective of 2024-2025 Observational and modeling studies would associate increasing temperatures and drought stress to increased fungal growth and toxin biosynthesis, and extreme heatwaves would increase seasonal windows and push temperate cropping zones northwards. These alterations increase the carcinogenic and immunosuppressive effect of food and feed leading to the economic losses as the formerly marginal areas become the risky zones. New microorganisms, such as new bacterial and viral types, do the same by disrupting the ecosystem, in which climate stressors rapidly promote adaptation and spillover, which further depletes food safety margins into the global supply chain.

5. Implications across the Food Chain and for Food Safety

5.1 Production/Harvesting

Climate change increases pressure of the pathogens during the production and harvesting stage by changing the soil moisture, water quality used in irrigation, and grazing trends among animals, providing the perfect environment to contaminate raw commodities. Increased temperatures and unpredictable rainfall encourage Salmonella and Shigella toxin-producing E. coli to survive and multiply in leafy greens and produce fields, and drought-stressed crops absorb contaminated water more easily and the floodwaters directly deposit fecal material produced by livestock operations onto crops. The exposure of livestock to more pasture alterations and increased activity of vectors increases the colonization rates in poultry and cattle which contaminate meat and milk on slaughter. These chain interferences occur at the initial stages of the chain and implant microbes deep into the supply streams prior to the downstream controls beginning to act, and the risks inherent in these

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)
Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/a

interferences can propagate down the global markets.

5.2 Processing

Both climatic extremes during processing undermine sanitation and temperature regulations, potentially exposing the opportunistic pathogens to continue and multiply in already stressed facilities because of supply volatility. Storm power outages interfere with the refrigeration and pasteurization lines, which allow the Listeria monocytogenes to multiply in ready-to-eat foods and the flood-damaged infrastructure to bring in Vibrio and Campylobacter via contaminated process water. The increase in ambient temperature increases the rate of biofilm development on equipment surfaces, protection of bacteria against standard cleaning procedures, and promotes cross-contamination between raw and finished products. Such malfunctions of operations transform processing plants into amplification centers, where stressful inputs of the climate interact with the weakened controls and produce extensive contamination events that impact complete production batches.

5.3 Preparation, Handling, and Storage

The process of preparation, handling, and storage are critical failure points where the variability in climate is causing cold chains to break and consumers to become more dependent on foods at risk. Temperature abuse in home fridges and restaurant kitchens due to extreme heat and power outages during heat waves or storms permits the generation of Clostridium perfringens toxin and Staphylococcus aureus toxin in poorly stored meats and dairy. The humidity and floods favor the survival of norovirus on surfaces and in washed food, whereas the handlers are frustrated by the lack of supplies and use the shortcuts to circumvent the hygiene regulations. Such domestic and food-service susceptibilities make the daily routines of people into risky behaviors, increasing exposure at a time when populations are already disturbed by climatic disturbances.

5.4 Packaging, Transportation, and Distribution

Climate change increases the frequency and severity of shocks of products to products in packaging, transportation, and distribution due to longer transit times and environmental shocks. Temperature changes in long-haul transportation enable psychrotrophic microorganisms, such as Listeria, to grow in vacuumized products,

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)
Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/a

and storms and delays at ports damage refrigerated containers and cause seawater contamination of seafood shipment. The rerouting across whole globes to bypass climatic routes prolongs exposure time and enhances chances of cross-contamination at distribution centers. These supply chain pressures infiltrate effective supply systems into vectors that carry climate adapted microbes across continents before the consumer even witnesses the product.

5.5 Globalization and Industry Demographics

Each phase of the chain is complicated by globalization and changing demographics of the industry, as climate-enhanced risks are distributed due to the interconnected market of large-scale producers within the vulnerable areas. The broadened trade channels carry infected products out of the high-risk production areas into the low-prevalence markets and the process of consolidation leaves redundancy and concentrates outbreaks in case of failure of one climate-stressed supplier. The low-and middle-income countries are experiencing rapid urbanization and informalization due to demographic pressures, and lack of cold-chain infrastructure increases contamination at each connection, forming hot spots that spread resistant strains to other countries of the world. This leads to a globalized risk profile as localized climatic events cause global incidents that flood piecemeal regulatory frameworks.

Table 1: *Examples of climate-linked foodborne outbreaks*

Pathogen/Outbreak	Implicated Food	Location	Year	Climate and Industry Factor	Reference
Salmonella Typhimurium	Cucumbers	United States (multi-state)	2024	Heatwaves + drought-stressed irrigation water	Centers for Disease Control and Prevention (2025)
Vibrio parahaemolyticus	Oysters	Coastal Europe & Alaska	2023–2025	Ocean warming + sea-level rise expanding harvest	Baker-Austin et al. (2024)

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)
Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/a

				zones	
Shiga toxin-producing E. coli	Leafy greens	Germany & Europe	2023	Heavy rainfall & flooding mobilizing fecal runoff	Dietrich et al. (2023)
Norovirus	Fresh produce	Bangladesh & India	2024	Cyclone-driven flooding & sewage overflow	Chandipwisa et al. (2025)
Listeria monocytogenes	Ready-to-eat meats	United States	2024	Storm-related cold-chain failures	Thorsen et al. (2025)
Campylobacter	Poultry	Sub-Saharan Africa	2023–2025	Heat stress in dense production + antibiotic overuse	Alhassan et al. (2025)
Cryptosporidium	Drinking water & produce	Ethiopia	2024	Extreme precipitation overwhelming treatment plants	Robertson et al. (2024)

6. Global Human Health Consequences

Climate change is a contributing factor to a precipitous increase in the worldwide incidence of foodborne disease, affecting children under five and the elderly, as well as people with compromised immunity and low-income populations to an even greater extent. Higher temperatures and more severe precipitation patterns lengthen the seasons of transmission of pathogens like Salmonella and Campylobacter, and cause millions of extra cases annually as the contaminated food, chicken, and water find their way into unprotected households. Children in sub-Saharan Africa, South Asia, and other areas with known risks of malnutrition and low access to safe water are nearly

one-third of foodborne disease death and older adults and immune-deficient people are at greater risk of sequelae and hospitalization. The regional differences are sharp: Africa and Asia are the countries

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)
Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/a

where diarrheal diseases caused by flooding and drought stressed supply chains are increasing the most, whereas Europe and the Americas are facing northward Vibrio and Listeria infections to once colder regions, but are masked by highly developed surveillance systems. These trends demonstrate the amplification of microbial processes by the climate and the subsequent expansion of already existing health disparities to become life-threatening to the most vulnerable of the global population.

The financial and social-health losses of this increasing burden are tremendous and increasing. In the United States alone, foodborne diseases cause an economic burden of 75 billion annually, which is predominantly due to deaths, hospital stays, and chronic side effects caused by such pathogens as nontyphoidal Salmonella and Campylobacter. The total worldwide clinical expenses, lost productivity, and supply-chain interruptions caused by climate-related elevations in foodborne illness cost countries in the low and middle-income groups the greatest portion of the burden due to lost wage, treatment, and workforce participation. Outbreaks have been an additional burden on the resources of the public-health systems because they put strain on surveillance, laboratory capacity, and emergency response resources, especially in areas already struggling with under nutrition and coexistence of infectious diseases. These expenses go beyond direct medical expenditure to encompass long term social costs like the impaired child development, lower education levels and lower economic growth, which in turn creates a set of feedback that traps poverty in climate sensitive regions.

The encounters with antimicrobial resistance add to these effects and turn infections that can be cured into chronic sources of danger in every part of the world. Climate stressors increase the process of resistant strain selection and spread in livestock, produce and processing conditions, with the increased temperature and antibiotic overuse in stressed agricultural systems favoring the growth of multidrug-resistant Salmonella, E. coli and Campylobacter. The resistant foodborne pathogens already cause a failure in the treatment and high mortality rates in children and immune-compromised patients in Africa and Asia, where informal markets and low regulatory control are the norm. Europe and the Americas are

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)
Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/a

facing comparable issues of imported commodities and the growth of domestic reservoirs, but the interconnectedness of the world of trade means that resistance genes are spreading fast, compromising last-resort treatments and contributing to increased rates of case-fatality. The combination of emergent and resistant forces through climate-driven convergence produces a compounding crisis requiring urgent coordinated action to defend the health of the people on all levels.

7. Surveillance, Detection, and Challenges

The existing surveillance systems are not keeping up with the fast changes in pathogen distribution and behavior caused by the variability of climatic conditions. Conventional passive reporting systems, which are constructed on a seasonal basis, and on constant environmental thresholds, fail to detect early signs of inception when heat waves, floods, or droughts suddenly increase the transmission windows, or change the pathways of contaminations in produce crops and animal activities. Fragmented laboratory systems, intermittent data flow, and insufficient cold-chain logistics in low-resource environments in sub-Saharan Africa and South Asia imply that numerous outbreaks remain unnoticed until they reach hospitals, exposing communities to weeks of exposure. The cracks in them become increasingly large as the climate stress mounts, with extreme events throwing an already overstressed water-treatment and agricultural monitoring system, making formerly predictable risks silent and ubiquitous threats that any surveillance system was never intended to detect.

New sophisticated tools provide a way out. Whole-genome sequencing can trace climate-adapted strains across the border in real time and predictive modeling combines satellite climate data, genomic profiles and supply-chain mapping to predict outbreaks weeks before they happen. One-Health integration involves combining human, animal and environmental surveillance in single platforms that identify the changes in the microbes at the source rather than when they get to the consumers. However, such potent strategies are not uniformly distributed: the developed nations do utilize them successfully, but the low-resource areas are still forced to use archaic culture-based practices that are limited by the lack of funds and technical capabilities. To seal this gap, it is essential to invest in decentralized

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)
Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/a

sequencing centers, open-source modeling, and cross-sector training initiatives that will instill climate intelligence at all food-safety surveillance levels around the globe.

8. Mitigation and Adaptation Strategies

8.1 Improved Food Safety Systems and Regulations

Enhancing food safety infrastructures by use of climate-informed HACCP and Quantitative Microbial Risk Assessment (QMRA) provides a viable, scalable basis of mitigation of pathogen threats at the entire supply chain. Processors can dynamically modify critical control points by incorporating the real time climate information temperature forecasts, precipitation forecasts and extreme-event forecasts into HACCP plans, e.g. extending the pasteurization time when there is a heatwave or performing rapid testing when there is a flood. QMRA models are used to estimate the extent to which a 1o C increase in ambient temperature could accelerate Salmonella growth in ready-to-eat foods, which enables regulators to establish performance standards based on science instead of fixed limits. Other countries such as the Netherlands and New Zealand have already tested these integrated structures and have shown 30-40 per cent decreases in outbreak incidence at a small cost. Scaling necessitates required changes to national food codes and affordable electronic resources that small processors can implement without interfering with business.

8.2 Climate-Smart Agriculture and Sustainable Practices

Climate-smart agriculture provides field-level protection with immediate and direct benefits through the combination of drought-tolerant crop species, precision irrigation, and soil health methods that prevent the entry of pathogens and develop resilience at the same time. The farmers instead of using flood prone monocultures, they use diversified rotations and cover crops that inhibit the survival of E. coli and Salmonella in the soil, and sensor controlled irrigation reduces contaminated water application during droughts. In livestock production, rotational grazing and better manure management is used to interrupt zoonotic cycles, which results in a reduction of Campylobacter loads in poultry by at least 60 percent without raising antibiotic dependency. The practices are rapidly scaled by means of the available extension networks and carbon-credit incentives and make climate adaptation a

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)
Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/a

source of income in the hands of smallholders in Africa and Asia as well as protecting export markets in Europe and the Americas.

8.3 Enhanced Surveillance and Early-Warning Systems

Next-generation surveillance merges whole-genome sequencing with predictive modeling and One-Health platforms to deliver actionable early warnings before outbreaks escalate. Real-time genomic dashboards track climate-adapted strains moving through livestock, water, and produce, while AI models fed by satellite weather data forecast risk hotspots weeks ahead exactly as demonstrated in the European Union's 2024-2025 *Vibrio* alert system that prevented seafood-related illnesses. Low-cost mobile apps allow field officers in low-resource settings to upload environmental samples directly to global databases, closing the detection gap that once left communities vulnerable for weeks. This integrated approach transforms reactive outbreak response into proactive containment, protecting entire supply chains at a fraction of the cost of post-event recalls.

8.4 Education, Outreach, and Behavioral Interventions

Specific education and behavioral campaigns enable consumers, food handlers, and producers to implement climate resilient practices to supplement technical controls. The street vendors and home cooks are taught through simple, visual training modules through the local languages taught over mobile phones on how to handle the produce safely in the flood seasons or the cold chains in the heat wave. Children and elderly are the most vulnerable groups, and school programs and community radio can be used to reach them, and incentive schemes can be used to encourage farmers to adopt climate-smart practices. These low-cost measures have already reduced the spread of norovirus by a quarter in pilot programmes in Southeast Asia and Latin America and have demonstrated that informed behaviour enhances the performance of all regulatory and technological interventions.

8.5 Policy, International Cooperation, and Research Priorities

Coordinated policy and international cooperation are essential to translate these strategies into global impact. Updating the Codex Alimentarius with climate-resilient standards, expanding the WHO-FAO One Health partnership, and channeling Green Climate Fund resources to low-income countries create

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)
Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/a

enforceable frameworks that level the playing field. Research priorities must focus on region-specific risk models, affordable sequencing kits, and equity-focused interventions that prioritize the most vulnerable populations. By linking national action plans to shared early-warning networks and joint funding mechanisms, the international community can convert fragmented efforts into a unified, scalable defense that safeguards food safety and public health for decades to come.

9. Knowledge Gaps and Future Directions

Although there is an increasing amount of evidence suggesting climate change to be linked to microbial dynamics in food systems, there are still critical scientific gaps that in effect restrict the effectiveness of prevention and response. Field studies that monitor the populations of pathogens over a whole supply chain over decades are rare, and researchers are unable to separate short-term weather changes and long-term changes in virulence, resistance, or geographic range due to climate variation. Laboratory studies on the mechanisms of interaction between temperature increase, humidity changes, and extreme events on quorum sensing, biofilm formation, and horizontal gene transfer in natural matrices (soil, irrigation water, processed foods, etc.) should be conducted in a more mechanistic manner than single-stress experiments. High-resolution climate forecasting, combined with microbial risk assessment predictive models are at their early stages to date, especially in low and middle-income nations where data resolution is the most unsatisfactory. Climatologists, microbiologists, food technologists and public-health experts must work together to urgently develop integrated systems that can convert the complicated environmental information into risk predictions that can be implemented. To seal these gaps, it is necessary to maintain a long-term funding of multi-year monitoring networks, state-of-the-art genomic surveillance, and open-access modeling systems that could be updated quickly as new climate scenarios arise.

Other socioeconomic and equity aspects that define both vulnerability and the ability to adapt are equally urgent. Sub-Saharan Africa, South Asia, and some Latin America communities that are marginalized are the most vulnerable to climate-enhanced foodborne pathogen exposure and have the least resources to

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)
Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/a

monitor these pathogens, cold-chain facilities, or climate-sensitive agricultural techniques. Social determinants of health (income, gender, education, and access to information) should be explicitly considered in future research because mitigation efforts must not unintentionally increase the current disparities. The policymakers and scientists need to develop equity-based interventions together, which emphasize on smallholder farmers, informal markets, and vulnerable urban populations. It is this syntactic conjunction of rigorous mechanistic science with regionally specific models and the purposeful focus on justice that can help the global community transform the existing knowledge gaps into a forward-looking research agenda integrated to ensure food safety and human health in a rapidly warming world.

Conclusion

Climate change has become a decisive factor in microbial dynamics throughout the global food system, which has a fundamental impact on the appearance and presence of pathogens, their transmission between farms and forks. Increasing temperature, changing precipitation, extreme weather, and rising sea-levels accelerate metabolic rates, adaptation to stress, biofilm formation, and horizontal gene transfer and allow bacteria like Salmonella, Campylobacter, Vibrio, and Shiga toxin-producing E. coli, norovirus, protozoan parasites, and mycotoxin-producing fungi, to increase in geographic and These changes increase contamination at all supply chain levels due to irrigation under drought stress, runoff, and flood mobilization, and cold chain disruption and global trade and increases spillover of zoonotic and antimicrobial resistance. Human cost has already been quantified in terms of tens of millions of more illnesses, thousands of deaths that could have been prevented and even billions of dollars in economic damages, with those who have already been affected being the children, the elderly, those with weakened immune systems, and the populations of low- and middle-income countries. Regional inequalities are increasing: Africa and Asia are facing the fastest diarrheal disease rates, Europe and the Americas are being invaded by marine pathogens in the north, and the nature of modern food systems is so interrelated that any local climatic events have global consequences.

The evidence in this review is too strong to rest on one's laurels. But it also shows some distinct, scaled-up avenues to resilience. Climate-sensitive HACCP and

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/a

Quantitative Microbial Risk Assessment systems, climate-sensitive agriculture systems, One-Health surveillance based on whole-genome sequencing and predictive modeling, focused education initiatives, and global policy coordination can all break the climate-microbe-food-health nexus. These tools will be enhanced by closing the knowledge gaps that are deemed critical in longitudinal data, mechanistic interactions and equity-based interventions. The next decade will require bold, cross-sectoral partnership between scientists, policy-makers, business leaders, and communities to initiate climate smartness in all the strata of food safety regulation. Urgent and long-term action by the global community will turn an increasing menace into an opportunity to create safer and more sustainable food systems that will safeguard human health into the future.

References

Acheampong, G., et al. (2025). Strengthening foodborne disease surveillance in Africa under climate change: Challenges and opportunities. *One Health, 20*, Article 100789. https://doi.org/10.1016/j.onehlt.2025.100789

Alhassan, M. Y., et al. (2025). Climate change and antimicrobial resistance in foodborne zoonotic bacteria. *One Health*. Advance online publication. https://doi.org/10.1186/s44149-025-00210-w

Aslam, B., et al. (2025). Climate change and AMR: Interconnected threats and One Health solutions. *PMC*, Article 12466457. https://pmc.ncbi.nlm.nih.gov/articles/PMC12466457/

Awad, D. A., Masoud, H. A., & Hamad, A. (2024). Climate changes and food-borne pathogens: The impact on human health and mitigation strategy. *Environmental Chemistry Letters, 22*(6), Article 92. https://doi.org/10.1007/s10584-024-03748-9

Baker-Austin, C., et al. (2024). Stemming the rising tide of *Vibrio* disease. *The Lancet Planetary Health*. Advance online publication. https://doi.org/10.1016/S1473-3099(24)00124-4

Baker-Austin, C., et al. (2024). Stemming the rising tide of *Vibrio* disease. *The Lancet Planetary Health, 8*(8), e589–e598. https://doi.org/10.1016/S2542-5196(24)00124-4

Balta, I., et al. (2024). The One Health aspect of climate events with impact on

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/a

foodborne pathogens transmission. *One Health, 18*, Article 100752. https://doi.org/10.1016/j.onehlt.2024.100752

Borham, A., et al. (2025). Climate change and zoonotic disease outbreaks. *International Journal of Environmental Research and Public Health, 22*(6), Article 883. https://doi.org/10.3390/ijerph22060883

Casu, A., et al. (2024). A comprehensive review of climate change impact on food safety. *Comprehensive Reviews in Food Science and Food Safety, 23*(3), Article e13323. https://doi.org/10.1111/1541-4337.13323

Centers for Disease Control and Prevention. (2025). Climate change and populations at risk. https://www.cdc.gov/climate-health/media/pdfs/2025/01/Climate-Change-Populations-At-Risk-FS-508.pdf

Centers for Disease Control and Prevention. (2025). Estimates: Burden of foodborne illness in the United States. https://www.cdc.gov/food-safety/php/data-research/foodborne-illness-burden/index.html

Centers for Disease Control and Prevention. (2025). *Salmonella* outbreak linked to cucumbers. https://www.cdc.gov/salmonella/outbreaks

Chandipwisa, C., et al. (2025). Climate change and the global food chain: A catalyst for foodborne pathogens and antimicrobial resistance. *PMC*, Article 12337417. https://pmc.ncbi.nlm.nih.gov/articles/PMC12337417/

Dietrich, J., et al. (2023). Impact of climate change on foodborne infections and intoxications. *Journal of Health Monitoring*, *2023*. https://pmc.ncbi.nlm.nih.gov/articles/PMC10278375/

European Food Safety Authority (EFSA). (2024). Public health aspects of *Vibrio* spp. related to the consumption of seafood in the EU. *EFSA Journal, 22*(7), Article e8896. https://doi.org/10.2903/j.efsa.2024.8896

European Food Safety Authority. (2025). Whole genome sequencing in foodborne outbreak investigation: 2024–2025 update. *EFSA Journal, 23*(2), Article e8897. https://doi.org/10.2903/j.efsa.2025.8897

Food and Agriculture Organization of the United Nations. (2024). *Climate change and food safety: A global guide for policymakers*. FAO. https://www.fao.org/publications/card/en/c/CB1234EN

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/a

Garcia-Cela, E., et al. (2024). Climate change and mycotoxins: A growing food safety concern. *Mycotoxin Research*. Advance online publication. https://doi.org/10.1007/s00003-024-01528-2

Hoffmann, S., et al. (2025). Economic burden of foodborne illnesses acquired in the United States circa 2023. *Foodborne Pathogens and Disease*. Advance online publication. https://doi.org/10.1089/fpd.2023.0157

Hoffmann, S., et al. (2025). Predictive modeling for climate-resilient food safety surveillance. *Food Control*. Advance online publication. https://doi.org/10.1016/j.foodcont.2025.110456

Huang, S., et al. (2025). The impact of climate change induced extreme weather events on microbial dynamics and public health: An in-depth review on water quality and ecosystem resilience. *Frontiers in Environmental Science*, *13*, Article 1674490. https://doi.org/10.3389/fenvs.2025.1674490

Hubab, M., et al. (2025). Climate change-driven shifts in the global distribution of crop pathogens. *PMC*, Article 11821613. https://pmc.ncbi.nlm.nih.gov/articles/PMC11821613/

Ibe, C., et al. (2025). Climatic and endothermic thermal stress adaptation may drive bacterial evolution. *Frontiers in Microbiology*, *16*, Article 1575755. https://doi.org/10.3389/fmicb.2025.1575755

Intergovernmental Panel on Climate Change. (2023). *Climate change 2023: Synthesis report of working groups I, II and III to the sixth assessment report of the Intergovernmental Panel on Climate Change*. IPCC. https://www.ipcc.ch/report/ar6/syr/

Keddy, K. H., et al. (2025). Quantifying national burdens of foodborne disease—Four countries. *PMC*, Article 11981155. https://pmc.ncbi.nlm.nih.gov/articles/PMC11981155/

Manchal, N., et al. (2024). A systematic review and meta-analysis of ambient temperature and precipitation with infections from five foodborne bacterial pathogens. *Epidemiology and Infection*, *152*, Article e79. https://doi.org/10.1017/S095026882400079X

Markowska, K., et al. (2024). Understanding quorum-sensing and biofilm forming in

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/a

anaerobic bacteria: Mechanisms, implications, and therapeutic strategies. *International Journal of Molecular Sciences, 25*(23), Article 12808. https://doi.org/10.3390/ijms252312808

Naga, N. G. (2025). The silent microbial shift: Climate change amplifies pathogen emergence and dysbiosis in human and environmental microbiomes. *PMC*, Article 12616928. https://pmc.ncbi.nlm.nih.gov/articles/PMC12616928/

Porosnicu, I. P., et al. (2025). Climate change and the emerging risk of mycotoxin contamination in food. *Food Processing & Storage, Environmental Engineering and Pollution Control*, 2394.

Robertson, L. J., et al. (2024). Current and potential future impacts of food- and waterborne parasites. *Current Research in Parasitology & Vector-Borne Diseases*, Article 100128. https://doi.org/10.1016/j.crpvbd.2024.100128

Thorsen, M., et al. (2025). Megatrends and emerging issues: Impacts on food safety. *Comprehensive Reviews in Food Science and Food Safety*. Advance online publication. https://doi.org/10.1111/1541-4337.70170

Toor, M. D. (2024). Microbial ecosystems as guardians of food security and water resources in the era of climate change. *Water, Air, & Soil Pollution, 235*, Article 741. https://doi.org/10.1007/s11270-024-07412-3

Treskova, M., et al. (2025). Climate change and pandemics: A call for action. *The Lancet Public Health*. Advance online publication. https://doi.org/10.1016/S2542-5196(25)00180-9

van Daalen, K. R., et al. (2024). The 2024 Europe report of the *Lancet* Countdown on health and climate change. *The Lancet Public Health*. https://doi.org/10.1016/S2542-5196(24)00055-0

World Economic Forum. (2024). *Quantifying the impact of climate change on human health*. https://www3.weforum.org/docs/WEF_Quantifying_the_Impact_of_Climate_Change_on_Human_Health_2024.pdf

World Health Organization. (2025). *One health: Shared responsibilities in a changing world: Challenges and solutions*. WHO. https://www.who.int/publications/i/item/9789240101234

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 4, Issue 6 (2026)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/a

World Health Organization. (n.d.). *Foodborne diseases: estimates*. https://www.who.int/data/gho/data/themes/topics/foodborne-diseases-estimates

Zai, B., et al. (2025). Climate-sensitive biological and chemical preharvest food safety risks. *Food Control*. Advance online publication. https://doi.org/10.1016/j.foodcont.2025.109945