Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

RECENT ADVANCES IN QUANTITATIVE GENETICS OF EUCALYPTUS: IMPLICATIONS FOR BREEDING AND CONSERVATION

Isha Shakoor
Institute of Botany, University of Punjab, Lahore,Pakistan.
ishashakoor07@gmail.com

Muhammad Umar (Corresponding Author)
Department of Agronomy, University of Agriculture Faisalabad , Pakistan.
themuhammadumar1@gmail.com

Shanzay Amjad
College of Statistical Sciences University of the Punjab, Pakistan.
shanzayamjad09@gmail.com

Atqa Rehna
Department of Botany University of Chakwal, Pakistan.
atqarehna77@gmail.com

Muhammad Ali Shah
Department of Chemistry, University of Chakwal, Pakistan.
alishah13382@gmail.com

Jamal Khabab Ahmad
Institute of Horticultural Sciences, University of Agriculture Faisalabad
j.Khabab642@gmail.com

Naeem Iqbal
School of Mathematics , Minhaj university Lahore, Pakistan.
nmathematics101@gmail.com

Nabila Arshad
Department of Chemistry, Mirpur University of Science and technology (MUST) Pakistan.
dr.nabila99999@gmail.com

ABSTRACT

Eucalyptus quantitative genetics research has shown remarkable growth but faces ongoing complications due to multi-trait optimization and genotype-environment

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

interactions along with natural and cultivated population diversity conservation requirements. The increasing need for sustainable forestry combined with upcoming climate change challenges requires precise and innovative breeding approaches that confront these obstacles. The integration of high-throughput sequencing with genomic selection (GS) and marker-assisted selection (MAS) help accelerate genetic gains and enhance climate resilience in Eucalyptus breeding programs according to this review. Scientists aim to close Eucalyptus' current performance-based gaps between its genetic potential and commercial and ecological requirements by applying new genomic technology platforms and quantitative genetic methods. This review shows how maintaining diversity among breeding targets requires scientists to prioritize wood quality traits and disease resistance together with drought tolerances. The advance of high-throughput sequencing methods and their application in SNP array development (EUChip60K) along with GWAS for complex traits and extended use of transcriptomics to understand stress responses represent major breakthroughs. The accuracy rate for GS predictions exceeds 0.7 for lignin content assessment and allows shorter breeding cycles and MAS helps discover QTLs for disease resistance along with wood quality. Genomic research has helped conservation scientists locate adaptive genetic markers which guide them in developing approaches to safeguard population genetic diversity under climate stress. Research findings demonstrate how various multi-omics datasets when paired with advanced phenotyping approaches together with AI-based predictive tools boost precision breeding and conservation success rates. Through the application of emerging technologies and the resolution of existing challenges quantitative genetics will preserve Eucalyptus as a fundamental element of sustainable forestry along with ecological resilience during global environmental transformations.

Keywords: Breeding Programs, Climate Adaptation, Genomic Selection, Quantitative Genetics, Sustainable Forestry

INTRODUCTION

The diverse genus of Eucalyptus featuring over 700 species maintains vital ecological and industrial values on a worldwide scale and creates significant economic opportunities. The native Eucalyptus species from Australia and its adjacent territories now thrive throughout tropical and subtropical regions because of its rapid growth speed alongside its adaptation ability and practicality which supports agricultural and forestry systems (Malakar, 2024). The use of Eucalyptus extends through wood materials to paper production and fragrance extracts to greenhouse gas storage and renewable energy generation thus playing an essential function in building sustainable systems. Renewed market interest in Eucalyptus products combined with environmental stressors including climate change and disease and scarcity of pests require new approaches to sustainably increase its productivity along with developing stronger resistance mechanisms (Ikeh, 2023; Varghese et al., 2024). The study of quantitative genetics assumed a central position as researchers aim to understand key trait heritance patterns while developing sustainable forestry breeding strategies (Gautam et al., 2024).

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Quantitative genetics through systematic investigation of multiple-gene inheritance patterns and environmental influences now guides Eucalyptus breeding practices (Li et al., 2024). Breeding techniques using phenotypic selection as their foundation face major restrictions because they deliver limited accuracy and require extensive breeding times. New statistical modeling approaches alongside molecular tools and genomic technologies have transformed the field by offering advanced methods for trait manipulation including growth rates and wood quality assessment and drought tolerance and disease resistance control (McGowan et al., 2021; Baldi & Porta, 2022). Gambling methodologies such as genomic selection alongside quantitative trait loci (QTL) mapping provide modern solutions for expedited genetic merit prediction (Jones, 2024). Current knowledge of trait heritability and genetic diversity together with genotype-by-environment interactions on trait expression remains limited because of ongoing gaps in our understanding. The effective sustainability of Eucalyptus resources depends on eradicating these essential knowledge gaps according to Climent et al. (2024).

Eucalyptus makes substantial economic contributions but its ecological conservation power alongside its potential in climate change mitigation remains equally important. Eucalyptus plantations operate as carbon storages and simultaneously benefit soil structure and maintain eco-system biodiversity (Hutapea et al., 2023). The long-term preservation of Eucalyptus ecological benefits depends on maintaining both genetic diversity within and between different Eucalyptus populations. Fundamental to adaptive potential is the preservation of genetic diversity which maintains successful outcomes in breeding programs (Bragg et al., 2021). Modern genomic tools including single nucleotide polymorphism (SNP) arrays and high-throughput sequencing enable our enhanced capacity for genetic diversity evaluation as well as conservation (Udoh et al., 2021). The broad implementation of genomic data in breeding pipelines meets resistance from two major challenges - the high expenses of genomic studies along with difficulties in genomic data integration. Scientists must now understand better how climate change affects the intricate relationship between genetics and environment which shapes adaptive traits according to Sinha et al., 2023.

This review examines the latest developments in Eucalyptus quantitative genetics while assessing their value for breeders and conservation programs. According to Ahmar et al. 2021 the paper explains three fundamental quantitative genetic research approaches which include family-based analyses QTL mapping and genomic selection. Quantitative genetic approaches enable scientists to measure trait heritability while discovering genetic markers which link to essential characteristics and build improved prediction models for breeding. The review examines Eucalyptus species genetic diversity within populations alongside diversity between species including its relevance for breeding and conservation practices (Mieres et al., 2024). A section follows the document which delves into the heritability properties of vital characteristics such as growth and wood quality alongside disease resistance and drought tolerance while explaining their role in breeding approaches. Review by Hashmi et al. (2023) examines advanced genomic tools and their utilization for marker-assisted selection together with high-throughput phenotyping applications to demonstrate their future possibilities in Eucalyptus breeding operations.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

This review focuses on studying the practical deployment of quantitative genetic research results towards breeding initiatives and wildlife conservation strategies. The paper presents breeding strategies to develop productive Eucalyptus varieties which exhibit disease tolerance and adaptive traits for changing climates together with conservation strategies to safeguard natural Eucalyptus genetic diversity (Bradbury et al., 2021; Getachew et al., 2024). The analysis examines ongoing challenges including difficult multi-trait evaluation processes and ensures the availability of durable genomic tools and considers how climate change influences breeding goals. The review makes clear recommendations for upcoming research by identifying present barriers that necessitate enhanced genomic methodology applications alongside conventional breeding practices according to Chaudhary et al. (2024).

This review demonstrates how quantitative genetics maintains essential value for Eucalyptus research and application advancement. Significant progress emerged from united traditional and modern approaches yet many essential steps remain to optimize the potential of the field completely. This review combines existing scientific literature with gap identification to suggestions for future work thus establishing foundational research paths while showcasing quantitative genetics potential for forest sustainability and biodiversity protection. This research gathers insights from methodological explorations and application examples alongside research outcomes to create a detailed view of quantitative genetic Eucalyptus research advantages and barriers for long-term improvement and conservation efforts.

METHODS OF QUANTITATIVE GENETIC ANALYSIS

Quantitative genetics research of Eucalyptus depends on multiple strong analytical methods to study genetic variations and calculate heritability estimates and forecast breeding success. The field adopts four primary methodologies which include family-based studies together with quantitative trait loci mapping and genomic selection and statistical approaches according to Müller et al. (2017). Current Eucalyptus genetics research benefits from these scientific methods as they provide breeding and conservation strategies with direct-targeted solutions.

Family-Based Studies

Family-based studies function as fundamental building blocks for quantitative genetic analyses. Scientists analyze genetic information in related individuals to obtain heritability estimates and separate total genetic variation into its additive and dominance and environmental elements (Serpico et al., 2023). In Eucalyptus research laboratories Progeny trials and half-sib or full-sib family designs serve as the principal analytical tools. Relevant research findings using this approach enabled scientists to discover important heritable elements such as growth speed alongside wood density resistance capability that sustain breeding initiatives together with conservation work (Mieres et al., 2024). Researchers measure narrow-sense heritability for wood quality traits through experiments using half-sib family designs. The predictive breeding values together with genotype selection for industrial use become possible through these research designs (Jones et al., 2021). Research based on family studies requires extensive resources which involve long periods of time and large expanses of land to create and sustain trial conditions. Family-based designs represent vital genetic data generation approaches despite extensive resource requirements (Lyu et al., 2024).

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

QTL Mapping

Eucalyptus genetics benefits substantially from the vital DNA mapping technique called Quantitative Trait Loci (QTL). Genomic segments called QTLs link to particular observable characteristics in populations. Researchers identify statistical connections between genetic markers and population-based phenotypic alterations through loci mapping processes (Zhu et al., 2023). Researchers extensively use both interval mapping and composite interval mapping techniques within their work. Researchers have used QTL mapping technology to discover genetic loci that control drought tolerance and disease resistance alongside lignin content in Eucalyptus trees. Several research works identified QTLs that regulate growth-related characteristics leading to important genetic insight about key economic traits (Du et al., 2022). High-density single nucleotide polymorphism (SNP) markers have integrated into QTL mapping procedures to enhance genetic variation detection precision at smaller scales.

QTL Mapping Application in Eucalyptus	Key Findings
Growth Traits	Identification of loci affecting height and biomass production.
Disease Resistance	Discovery of QTLs linked to rust and fungal resistance.
Wood Properties	Genetic loci associated with cellulose and lignin content.

Genomic Selection

Genomic selection transforms Eucalyptus breeding by using whole-genome markers to forecast the genetic properties of various individuals. The approach differs from classic studies that target targeted loci because genomic selection evaluates every available marker while detecting variations with both minor and major impact (Wang & Zhang, 2024). Widespread acceptance of this method emerged because of its enhanced selection accuracy and need to shorten breeding lifecycles. Genomic selection uses high-throughput sequencing technology to deliver highly precise forecasts of traits including growth rate and wood quality and drought tolerance levels (Persa et al., 2021). Genomic selection offers significant potential for Eucalyptus through accelerated experimentation which frees resources for optimized deployment (Duarte et al., 2024). The process of genomic selection demonstrated its ability to enhance genetic gains for resistance-based traits. Scientists observe enhanced forecasting success by including environmental factors into mathematical models which consider genotype-by-environment interactions. Genomic selection has proved its ability to examine challenging traits while demonstrating both flexible characteristics and strong operational capabilities (Anilkumar et al., 2022).

Statistical Methods

Quantitative genetic data analysis requires statistical methods which offer essential tools for precise estimation together with prediction capabilities. Within Eucalyptus genetics research Mixed linear models (MLMs) together with restricted maximum likelihood (REML) stand as the primary analytical approaches.

- **Mixed Linear Models (MLMs):** Phenotypic variance partitioning happens through these models which combine fixed and random effects while separating genetic and environmental components. The tools prove essential for QTL

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

mapping and genomic selection studies because they permit researchers to operate on complex data structures (Misztal et al., 2021).

- **Restricted Maximum Likelihood (REML):** Mixed Linear Models (MLMs) use REML techniques to produce highly precise estimates of components that explain variance between plant populations. Family-based studies heavily benefit from this method because precise variance partitioning forms the core requirement for heritability estimation (Tang et al., 2022).

The application of these statistical methods ensures robust analysis of genetic data, facilitating the identification of key traits and informing breeding decisions. The combination of advanced statistical models with genomic tools has greatly enhanced the predictive power and efficiency of quantitative genetic analysis in Eucalyptus (Alemu et al., 2024).

Statistical Method	Application	Advantages
Mixed Linear Models	QTL mapping, genomic selection	Handles complex datasets, accounts for random effects.
REML	Variance component estimation in family-based studies	High precision in partitioning variance.

Quantitative genetics studies in Eucalyptus now use family-based investigations in combination with QTL mapping together with genomic selection and statistical models. These methods create detailed examinations of heritability and genetic variation and thereby enable advanced targeted breeding practices (Dhakal et al., 2023). Research using advanced genomic tools combined with statistical models lets scientists address fundamental Eucalyptus breeding and conservation challenges to produce suitable outcomes for this essential group of plants (Jordan et al., 2023).

GENETIC DIVERSITY IN EUCALYPTUS SPECIES

Diversity Within and Between Species

More than 700 species comprise the Eucalyptus genus which shows outstanding genetic and phenotypic diversity between its members. The extensive genetic variety enables Eucalyptus' flexible response to different environments and supports its applications across industries which produce pulp and paper and generate bioenergy as well as provide essential oils (Kim, 2024). The extensive analysis of Eucalyptus species variations helps scientists track evolutionary history and develop breeding and conservation initiatives (Pfeilsticker et al., 2022).

Wide-ranging species Eucalyptus globulus and Eucalyptus camaldulensis exhibit strong intra-specific genetic diversity between their populations. Population-based studies show significant distinctions in growth attributes and wood density as well as drought adaptation and disease immunity throughout the ranges of these species (Malakar, 2024). Different communities of E. globulus across Australian regions show distinct wood quality along with unique growth characteristics due to their local adaptation to variable climates and soil conditions (Callister et al., 2024).

Moreover inter-specific diversity provides genetic developers with potential to enhance their population by merging important features originating from different species. Hybrids created from Eucalyptus grandis and Eucalyptus urophylla show widespread

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

commercial use in forestry because they thrive in tropical environments and display rapid growth rates according to Lee et al. (2023). Inter-specific diversity demonstrates that this plant genus can be advanced through directed breeding initiatives because of its practical applications.

Impact of Genetic Diversity on Breeding and Conservation Programs

Genetic diversity functions as the critical foundation for successful breeding combined with conservation methods in Eucalyptus species. Breeding programs achieve their greatest benefits when they maintain extensive genetic variation that lets breeders choose optimal characteristics for improved productivity alongside stronger adaptive capabilities. Scientists have implemented breeding programs successfully in Eucalyptus plantations by applying genetic diversity approaches to develop fast-growing timber with high wood density along with enhanced pest protection (Thapliyal et al., 2023). Recurrent selection stands out as an important breeding program because professionals select outstanding individuals across successive generations to improve traits. Application of this method has achieved better growth rates and frost resistance when used with Eucalyptus nitens plants. These programs succeed only when genetic diversity gets maintained because inbreeding depression leads to decreased fitness and diminished adaptability (Rocha 2023).

The long-term survival of many Eucalyptus species depends on maintaining genetic diversity to protect them from current threats of habitat loss and climate change and pest attacks. Protection strategies including in situ reserves and ex situ seed banks seek to maintain full genetic diversity representation for widespread as well as uncommon Eucalyptus species (Negash, 2021). Conservation programs targeting the endangered Eucalyptus morrisbyi seek to protect multiple genetic variants that create its capacity to withstand climate modifications.

Advances in Genomic Tools for Mapping Genetic Variation

Genomic tools have transformed Eucalyptus genetic diversity assessment by allowing scientists to undertake precise mapping of genetic divergence. High-throughput sequencing technologies alongside genomic resources including single nucleotide polymorphism (SNP) arrays and whole-genome sequencing reveals essential information about Eucalyptus key trait genetics (Trudić et al., 2024; Gautam et al., 2024). Genetic analysis across different populations as well as species obtains significant assistance from SNP array technology. Genetic markers linked to growth traits together with wood quality and disease resistance have been identified through SNP genotyping research. The markers allow marker-assisted selection (MAS) to advance breeding programs through their ability to detect desired traits in early developmental stages (Ahmar et al., 2021; Mori & Cipriani, 2023).

Whole-genome sequencing based technology has led to enhanced comprehension of genetic variations throughout Eucalyptus populations. The Eucalyptus grandis genome sequence established an important framework which supports comparative analysis throughout the Eucalyptus genus (Ferguson et al., 2024; Li et al., 2024). Genomic studies identified highly important genes active in lignin biosynthesis and cellulose production and secondary metabolite pathways for industrial applications (Wang et al., 2021).

The exploration of genetic variation and phenotypic traits in Eucalyptus became

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

possible through extensive use of Genome-wide association studies (GWAS). Genetic factors linked to pulp yield and wood density in E. globulus were exposed through genome-wide association studies leading to potential targets for breeding improvement (Valenzuela et al., 2021; Rocha et al., 2022). Genomic tools demonstrate their potential to make breeding programs more precise and efficient according to studies conducted in this field.

Table: 3 Summary of Genetic Diversity and Genomic Tools in *Eucalyptus* Species

Aspect	Description	Key Studies/Examples	Applications
Intra-Specific Diversity	Variability within species, e.g., growth rate, wood density, drought tolerance.	*E. globulus* populations in Tasmania and Victoria show distinct wood quality traits (Butler et al., 2022).	Breeding programs for trait-specific improvements (e.g., wood density, growth).
Inter-Specific Diversity	Variability between species, utilized in hybrid breeding programs.	*E. grandis* × *E. urophylla* hybrids used in tropical plantations (Hua et al., 2022).	Enhanced productivity and adaptability through hybrid vigor.
Role in Breeding Programs	Use of genetic diversity to improve economically important traits.	Recurrent selection in *E. nitens* for frost resistance (Tomé et al., 2021).	Improved plantation performance and resilience to environmental stress.
Role in Conservation Programs	Preservation of genetic variation in endangered species.	Conservation of *E. morrisbyi* genetic diversity for climate adaptability (Lott & Read, 2021).	Genetic diversity aids in maintaining species resilience and adaptability.
SNP Arrays	High-resolution markers for genetic variation mapping.	SNP genotyping of *E. globulus* identified loci for wood density (Valenzuela et al., 2021).	Marker-assisted selection for efficient breeding programs.
Whole-Genome Sequencing	Comprehensive mapping of genetic diversity and candidate genes.	Sequencing of *E. grandis* genome revealed lignin and cellulose biosynthesis pathways (Liu et al., 2021).	Identification of target genes for industrial applications.
GWAS	Linking genetic variation with phenotypic traits.	GWAS in *E. globulus* linked loci to pulp yield and wood density (Dasgupta et al., 2021).	Precision breeding through trait-specific genetic markers.
Climate Change Implications	Genetic diversity aids in adaptation to changing environments.	*Eucalyptus camaldulensis* populations show varying drought tolerance (Amrutha et al., 2021).	Breeding for resilience to climate variability.

This table summarizes key aspects of genetic diversity, relevant studies, and their applications in breeding and conservation programs.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about



A flowchart showing how genetic diversity is utilized in breeding programs, from selection to hybridization

Heritability of Key Traits in Eucalyptus

The genetic potential of evolutionary development within species relies on heritability to identify key traits for breeding programs that enhance ecological and economic characteristics (Shahrajabian et al., 2021). Estimates of heritability in the important timber genus Eucalyptus help identify genetic influences on essential traits including growth patterns and disease protection together with wood properties and drought resistance capability. A detailed evaluation explores how these important characteristics inherit over generations while examining their measurement approaches and the relevance to breeding operations according to Baldi & La Porta (2022).

Heritability of Growth-Related Traits

Economic sustainability of Eucalyptus plantations relies on three fundamental growth-related traits which consist of height projection and diameter at breast height (DBH) and volume development. Quantitative genetics research shows growth-related traits consistently demonstrate moderate to high heritabilities with values typically distributed between 0.3 and 0.6 across species and environmental situations (Mieres et al., 2024). Laboratory findings demonstrate that genetic control of DBH in Eucalyptus globulus reaches an estimated 0.5. Scientists calculate the selection potential of subjecting Eucalyptus grandis to breeding selection based on its heritability value range from 0.4 - 0.6 (Ibarra et al., 2023).

Field measurements of growth traits proceed in basic ways since they can be directly observed in their natural settings. The measurements of height rely on clinometers or laser rangefinders to determine the information and diameter breast height relies on

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

calipers or diameter tapes according to Duncanson et al. (2021). Measurement timing over extended periods results in continued data collection which enhances the precision of heritability calculations in successive time increments. Simple selection procedures based on phenotypic traits alongside family-based breeding programs prove successful for enhancing these traits because their heritability levels remain high. Soil quality together with water availability must be considered in breeding programs to achieve highest possible genetic improvement (Hasan et al., 2024).

Heritability of Disease Resistance

Eucalyptus breeding programs need disease resistance traits because pathogen outbreaks from changing climates and expanding plantations have become more common. Disease resistance heritability rates prove lower than those of growth-related traits with typical values between 0.1 and 0.3 (Hurel et al., 2021). Disease resistance tests for Mycosphaerella leaf disease in Eucalyptus globulus show around 0.2 heritability value but environmental factors still strongly influence disease expression (Rocha, 2023).

The evaluation of disease resistance requires inspecting symptom intensity through scored examinations or automated digital image collection methods. Assessments of disease symptoms take place under inoculation experiments and natural infection conditions using scoring systems that rate symptoms from no symptoms (0) to severe symptoms (5) (Laidig et al., 2021). Current advancements in phenotyping technologies through methods such as hyperspectral imaging and molecular diagnostics lead to improved measurement precision which enhances heritability assessment accuracy (Xiao et al., 2022). The low heritability of disease resistance necessitates the adoption of advanced breeding strategies. The breeding approach termed Marker-Assisted Selection (MAS) together with genomic selection (GS) addresses the deficit of low heritability through genetics-based breeding value prediction according to Yunus et al. (2025).

Measurement of disease resistance involves evaluating the severity of symptoms, often using scoring systems or digital imaging techniques. These assessments are typically conducted in controlled inoculation experiments or natural infection conditions, with scoring systems ranging from 0 (no symptoms) to 5 (severe symptoms) (Laidig et al., 2021). Advances in phenotyping technologies, such as hyperspectral imaging and molecular diagnostics, have improved the precision of these measurements, enabling more accurate heritability estimates (Xiao et al., 2022). The low heritability of disease resistance necessitates the adoption of advanced breeding strategies. Marker-assisted selection (MAS) and genomic selection (GS) have shown promise in overcoming the limitations of low heritability by leveraging genomic information to predict breeding values (Yunus et al., 2025). For example, genomic prediction models incorporating single-nucleotide polymorphism (SNP) markers have been successfully applied to identify disease-resistant genotypes in Eucalyptus species (Younessi & Gailing, 2022).

Heritability of Wood Quality Traits

The pulp and paper industry relies heavily on wood quality traits because density, fiber length, cellulose content and lignin composition shape product outcomes. The heritability levels of these traits fall into the medium-to-high range from 0.3 to 0.6 according to Rocha (2023).

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

The heritability estimates for wood density in Eucalyptus grandis come in at approximately 0.4 whereas the estimates for cellulose content in Eucalyptus globulus reach up to 0.5 (Rocha, 2023; Ibarra et al., 2023). Ethical measurement of wood traits involves the use of specific technology protocols. The evaluation of density and stiffness traits in materials now mainly relies on non-invasive methods such as near-infrared (NIR) spectroscopy combined with acoustic velocity testing. The analytical process of destructive sampling combines core extraction with laboratory-based chemical analysis methods to extensively measure fiber properties and chemical composition (Olaoye & Ojo, 2022).

Heritability of Drought Tolerance

Eucalyptus species need strong drought tolerance capabilities because they exist in arid and semiarid regions. Drought tolerance heritability studies demonstrate that genetic influence on these traits ranges between 0.2 and 0.4 based on estimates from González (2023). A wide range of heritability values exists between drought tolerance characteristics including water-use efficiency (WUE) and stomatal conductance and root system architecture. WUE measurements in Eucalyptus camaldulensis show heritability at 0.3 according to Lu & Fricke (2023) while root depth studies display a higher heritability value of 0.4. Drought tolerance assessment requires combined testing of biological traits along with observable features. Using porometers and pressure chambers researchers measure physiological traits including leaf gas exchange and osmotic potential. Morphological traits including root depth and shoot-to-root ratio are measured through destructive sampling and ground-penetrating radar according to Boursiac et al. (2022).

Research on drought tolerance indicates selection methodologies may produce modest genetic progress because of the trait's moderate heritability status. The biological advancement of drought-tolerant genotypes depends on genomic technologies that breeding programs now integrate into selection decisions (et al., 2022). Genome-wide association studies (GWAS) have detected genetic markers which help scientists identify drought tolerance genes within Eucalyptus providing new insights for marker-assisted breeding.

Implications for Breeding Programs

The successful development of breeding programs depends fundamentally on estimates of heritability in traits. Straightforward selection methods achieve genetic gains quickly for traits that demonstrate high heritability such as growth and wood quality. Conversely, traits with low heritability, such as disease resistance and drought tolerance, require more sophisticated approaches, including:

- **Marker-Assisted Selection (MAS):** Through molecular marker analysis scientists identify links between beneficial traits which boosts selection precision when working with traits that exhibit low heritability (Singh et al., 2021).
- **Genomic Selection (GS):** Through GS breeding scientists predict individual genetic worth by employing markers from across the entire genome to compensate for minimal phenotypic rating techniques (Anilkumar et al., 2023).

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

- **Hybrid Breeding:** Through hybridization of genetically diverse groups breeders can exploit the beneficial growth patterns of hybrid organisms to enhance weakly inherited traits including disease resistance (Begna, 2021).
- **Genetic Engineering:** Progress in genetic modification technologies including CRISPR-Cas9 provides scientists the ability to specifically modify vital trait genes beyond traditional breeding boundaries (Wang et al., 2023).
- **Environmental Interventions:** Park programs aiming to improve drought tolerance utilize environment enhancement techniques with enhanced irrigation methods and improved soil care to maximize genetic improvement (Xing & Wang, 2024).

Heritability estimates help breeding programs make decisions regarding trait selection priorities in Eucalyptus breeding. Breeding programs can achieve significant genetic improvements through classical selection for high heritability traits which include both growth values and wood quality features. Trace inheritance traits need integrated solutions with managed breeding technologies to achieve outcomes in disease resistance and drought tolerance together with environmental management systems. The implementation of these breeding methodologies enhances both the economic worth and ecological stability of this critical genus by improving productivity together with increased resilience.

Technological Advancements in Phenotyping and Breeding			
Technology	**Application**	**Example Traits**	**Advantages**
Near-Infrared (NIR) Spectroscopy	Non-destructive wood quality assessment	Wood density, cellulose	High throughput, precise measurements
Hyperspectral Imaging	Disease resistance and stress phenotyping	Pathogen infection severity	Accurate, real-time monitoring
Genomic Selection (GS)	Prediction of breeding values	Drought tolerance, disease	Accelerates breeding cycles
GWAS	Identification of trait-associated loci	All traits	Pinpoints genetic markers

Technological Advancements in Phenotyping and Breeding

Table 4: Summary of Heritability Estimates for Key Traits in Eucalyptus

Trait	Species	Heritability Estimate (h^2)	Measurement Method	References
Growth (e.g., DBH, height)	*E. globulus*	0.3–0.6	Field measurements (calipers, clinometers)	(Ribeiro et al., 2023)

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

Disease Resistance	*E. globulus*	0.1–0.3	Disease scoring (natural or artificial infection)	(Bartholomé et al., 2020)
Wood Density	*E. grandis*	0.4–0.6	NIR spectroscopy, core sampling	(Makouanzi et al., 2018)
Drought Tolerance	*E. camaldulensis*	0.2–0.4	Gas exchange analysis, root depth measurements	(Souden et al., 2020)

GENOMIC TOOLS IN EUCALYPTUS BREEDING

Eucalyptus represents a tree genus with 700 distinct species which serves as one of the world's most significant tree families from both a commercial and ecological perspective. The diverse industrial uses of Eucalyptus pulp continue to be fundamental to likely humanity's ability to pursue sustainable forestry practices. The traditional selection method of Eucalyptus breeding programs incorporating phenotypic characteristics proves effective but consumes significant resources and time (Zhao et al., 2022). Genomic technology has revolutionized Eucalyptus breeding procedures by improving efficiency while enabling precise measures and resistance-based and climate-related solutions. This section evaluates modern improvements in high-throughput sequencing alongside genomic selection (GS) and marker-assisted selection (MAS) applications during Eucalyptus breeding research (Feng et al., 2024).

High-throughput sequencing

High-throughput sequencing (HTS) platforms which comprise next-generation sequencing (NGS) systems have led to a genomics revolution through their power to analyze entire genomes swiftly yet economically. Modern advancements in sequencing technology have delivered essential genetic information about major Eucalyptus traits to breeding programs (Wang, 2024).

Genome assembly and annotation

Scientists published the entire genome sequence of Eucalyptus grandis in 2014 which established a new benchmark for Eucalyptus genomic research. The reference genome enables researchers to conduct comparative genetic studies which help identify genes affecting economically relevant traits including wood density and lignin content and disease resistance (Gupta & Shankar, 2024). Third-generation sequencing tools PacBio and Oxford Nanopore deliver improved genome alignment results which reveal phenotypic trait controlling insertions and deletions and duplications (Espinosa et al., 2024).

Applications in breeding

Modern sequencing technology enabled researchers to create dense single-nucleotide polymorphism (SNP) arrays like the EUChip60K SNP array which targets Eucalyptus species (Aguirre et al., 2024). Because these arrays provide technology for breeders to conduct genetic trait identification through genome-wide association studies (GWAS). GWAS analysis successfully associates specific SNPs with Eucalyptus growth attributes alongside wood attributes and Leptocybe invasa resistance (Calazans, 2023). Research on Eucalyptus species using RNA sequencing (RNA-seq) technology generated information about gene expressions at different environmental and biotic stress conditions (Pinheiro, 2021). The study of gene expression through

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

transcriptomics has shown that Puccinia psidii (myrtle rust) activates particular genes that led to the development of disease-resistant tree varieties (Manzoor et al., 2024).

Genomic Selection

Genomic selection (GS) marks a new direction in tree breeding through its ability to predict breeding values using comprehensive genome marker data. A major distinction exists between GS and standard selection procedures because GS relies on genomic data to increase its precision while diminishing breeding cycles.

Theoretical Basis and Implementation

Genetic selection (GS) calculates genomic estimated breeding values (GEBVs) from statistical models that incorporate genome-wide marker data to determine individual values. The predictive model includes key and minor influencing loci to deliver complete genetic measurement capabilities (Misztal et al., 2021). The implementation of GS for Eucalyptus breeding becomes achievable because of available dense SNP markers alongside robust estimation systems including genomic best linear unbiased prediction (GBLUP) and Bayesian methods (Sharma et al., 2024).

Benefits in Eucalyptus Breeding

The main benefit of applying GS technology allows breeders to cut down breeding cycle durations for Eucalyptus species which live very long. The lengthy juvenile phase together with phenotypic evaluation times normally stretches breeding cycles across many decades. The genetic screening method enables seed selection of superior genotypes which dramatically shortens the breeding timeline (Grattapaglia, 2022).

Research findings show that GS performs efficiently for forecasting growth rate together with wood quality characteristics and disease resistance within Eucalyptus species. Scientists demonstrated the practical application potential of GS breeding technology by achieving prediction accuracy rates above 0.7 for Eucalyptus urophylla's stem diameter and lignin content traits (Paludeto et al., 2021). GS demonstrates value in identifying low heritability traits including drought tolerance by utilizing genome-wide markers which discover small-effect loci (Jamil et al., 2024).

Marker-Assisted Selection (MAS)

The breeding process receives enhanced efficiency through the application of genetic markers that track traits of interest with marker-assisted selection (MAS). The genome-wide approach of GS contrasts with MAS which chooses marker loci that control major effect traits. Marker-assisted selection demonstrates its most potent impact on Eucalyptus breeding through its ability to handle traits that depend on key genes for resistance and wood composition (Sharma et al., 2024; Wang & Zhang, 2024).

Identification of Candidate Genes

Genomic tools developed recently have become effective for locating quantitative trait loci (QTLs) markers that show strong linkage to economically beneficial traits. The identification of loci linked to Cryphonectria cubensis (Eucalyptus canker) and Puccinia psidii (Kumar et al., 2024) resistance happened through QTL mapping. Resistant MAS varieties have incorporated these markers to their breeding programs.

Practical applications

The application of MAS technology has become prevalent throughout wood property improvement projects in Eucalyptus since cellulose and lignin represent essential factors within pulp and paper manufacturing. DNA markers linked to cellulose synthase

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

genes successfully enhance cellulose content during breeding population development (Connell, 2022). Researchers utilize MAS technology to enhance frost tolerance and rooting capabilities during clonal propagation operations. MAS delivers significant benefits in Eucalyptus breeding because it produces new hybrid varieties with improved pest and disease resistance. Breeders achieved the development of superior hybrid varieties through marker selection of resistance genes which perform well across diverse environmental settings (Younessi et al., 2022).

Genomic tools such as high-throughput sequencing and genomic selection and marker-assisted selection have dramatically transformed the methodology of Eucalyptus breeding. Through these technologies researchers have boosted the efficiency and precision of their breeding operations which produced new varieties with better growth characteristics combined with higher wood quality standards and disease fighting capabilities. Through improved management of genomic applications alongside emerging technologies Eucalyptus breeding programs can develop more resistant and productive strains for sustainable forests and conservation practices.

Table 5: Genomic tools, their applications, and recent advancements in Eucalyptus breeding:

Genomic Tool	**Key Features**	**Applications in Eucalyptus Breeding**	**Recent Advancements**
High-Throughput Sequencing	Enables rapid and cost-effective analysis of genomes	- Development of SNP arrays - GWAS for traits like growth, wood quality, and pest resistance - Transcriptomic studies for stress response	- Complete genome of *Eucalyptus grandis* - Improved assemblies using PacBio and Oxford Nanopore - EUChip60K SNP array - RNA-seq for gene expression studies
Genomic Selection (GS)	Predicts breeding values using genome-wide marker data	- Accelerates selection for traits like growth rate, wood quality, and disease resistance - Addresses complex traits like drought tolerance	- High prediction accuracy (>0.7) for traits like lignin content - Models like GBLUP and Bayesian methods - Applications in reducing breeding cycle time
Marker-Assisted Selection	Uses markers linked to specific traits to guide selection	- Targets major effect traits like disease resistance and wood quality - Enhances hybrid breeding programs	- Identification of QTLs for resistance to *Cryphonectria cubensis* and *Puccinia psidii* - Markers for cellulose and lignin traits - Integration into clonal propagation programs for hybrids
Challenges	Heterozygous genomes, large population sizes,	- Complexity in genome assembly and variant	- Pangenomics and single-cell sequencing emerging as

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

	and extensive gene flow	detection	solutions - AI and machine learning enhancing prediction accuracy for complex interactions
Future Directions	Integration of multi-omics data and advanced computational tools	- Holistic understanding of trait architecture - Precision breeding	- Leveraging metabolomics, phenomics, and environmental data - Machine learning for genomic prediction models - Greater incorporation of adaptive loci in breeding programs

RECENT ADVANCES IN QUANTITATIVE GENETICS OF EUCALYPTUS

Applications and Implications for Breeding and Conservation

Advanced breeding strategies empowered by quantitative genetic approaches now efficiently tackle three major plant production challenges which include disease resistance improvement and stress tolerance increases and yield optimization. Eucalyptus which ranks among the top planted genera worldwide for timber and bioenergy production requires advanced breeding strategies that achieve sustainable production together with environmental conservation of natural population genetic diversity (Healey et al., 2022).

Breeding for Productivity

Eucalyptus plantations achieve higher productivity because Quantitative genetic research has proven essential for trait discovery and selection. The advancement of quantitative trait loci (QTL) mapping combined with genome-wide association studies (GWAS) allows precise measurement of traits which include fast growth rates and precise determination of wood density and pulp yield (Mathiazhagan et al., 2021). Selection indices based on breeding programs use these tools as an acceleration method to attain notable genetic progress by quantifying crucial economic traits together within one selection system. Breeders leveraging high-throughput phenotyping systems alongside genomic selection programs now sustainably propagate future generations that demonstrate peak growth rates and optimal wood composition (Sandhu et al., 2022).

Hybrid Eucalyptus varieties have been developed through quantitative genetics principles that enable the utilization of heterosis effects in hybrids. Researchers using genetic variance components analysis can predict hybrid clone performance to deploy best-performing clones across diverse environments (Chaves et al., 2024).

Breeding for Disease Resistance

Worldwide Eucalyptus plantations sustain major economic damage from pathogen-caused plant diseases. Disease resistance traits have been deeply examined through quantitative genetic principles (Amaral, 2021). Disease resistance against fungal pathogens including Mycosphaerella spp. and rust (Puccinia psidii) has been measured

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

using pedigree-based along with genomic relationship models. Breeding through marker-assisted selection (MAS) and genomic selection (GS) has been made possible by identifying major resistance genes alongside polygenic contributions (Mapari & Mehandi, 2024). Resistance to Phytophthora cinnamomi pathogen has been integrated as a valuable addition in Eucalyptus varietals which prevent root rot. Through quantitative genetic approaches breeders can detect resistance in their breeding pool at early stages of development which reduces their need for extensive and costly field trials (Jeff et al., 2021).

Climate Adaptation

Eucalyptus serves as an excellent model for understanding climate adaptation research because it exists across a vast geographic area and diverse ecological habitats. Quantitative genetics reveals adaptive traits that involve drought tolerance along with frost resistance and heat stress resilience according to Prober et al. (2022). Research-based G×E examination makes possible the identification of genotypes that display robust performance throughout multiple climate zones.

Multiple physiological and morphological factors influence drought tolerance via their effects on water-use efficiency (WUE) and deep-rooting systems infrastructure (Sofi et al., 2021). The combination of quantitative trait mapping and genomic selection helped researchers understand important plant traits leading to the development of stress-tolerant varieties that can thrive in drought zones (Zampieri et al., 2023). Natural population studies of Eucalyptus adaptive genetic diversity guide current efforts to protect resilient gene pools for climate conservation.

Conservation of Genetic Diversity

Natural and managed Eucalyptus populations require conservation of genetic diversity because this preserves their long-term sustainability and improves resilience to environmental change. Natural population analyses performed by quantitative genetics reveal population-wide additive variation levels together with degrees of interbreeding and effective grain size dynamics (Jordan et al, 2023). These conserved metrics help determine the preservation strategies which protect genetic diversity including adaptation potential.

Conserving genetic diversity becomes essential for Eucalyptus regnans and Eucalyptus grandis species because they endure habitat fragmentation along with environmental range shifts caused by climate change. The use of quantitative genetic information allows conservationists to create seed collection approaches that normalize germplasm storage methods to shield comprehensive genetic diversity (Fahey, 2023; Kulak et al., 2022). Gene flow assistance supported by genetic data enhances the adaptive abilities of populations that need to survive changing environmental conditions.

Challenges and Future Directions

Important progress has been achieved in quantitative genetics studies of Eucalyptus research but several persistent hurdles restrict its complete exploitation as a breeding tool and biodiversity management solution.

Complexity of Multi-Trait and Genotype-by-Environment Interactions

Programs that conduct Eucalyptus breeding seek to enhance several traits at once particularly growth together with wood quality attributes and disease resistance characteristics. The optimization of single traits tends to reduce the possible

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

improvement of related traits according to Silva et al. (2024). The quest for faster growth in selections could trigger unintended changes which lead to decreased wood density. Multi-trait models featuring genetic correlations and G×E interactions become essential for grasping and controlling these trade-offs (Fradgley et al., 2023).

Research efforts aimed at breeding face additional obstacles due to genotype-by-environment interactions which produce dissimilar performance levels between genotypes across diverse environments. The intricacy of G×E effects requires both advanced statistical mapping and experimental methods to measure them accurately. Long-term multi-environment trials and genomic prediction work together to uncover stable adaptable genotypes (McBreen et al., 2025).

Need for Robust Genomic Resources

The genomic resources available for Eucalyptus, while substantial, still lag behind those of other major crops. The accuracy of quantitative genetic evaluations depends on high-quality reference genomes together with annotated transcriptomes (Wang et al., 2023). New dense SNP array technologies together with pangenomic methods will improve genetic variation assessment and help identify previously unknown alleles (Ogunbawo et al., 2024). Genotyping and phenotyping operations present financial constraints that especially affect limited breeding operations because of their restricted resources. Scientific improvements in sequence analysis methods together with testing systems will be essential to remove these blocking factors.

Role of Climate Change

The current climate crisis creates unique threats for both Eucalyptus breeding operations and natural conservation projects. Eucalyptus breeding requires the adoption of resilient varieties because temperatures keep rising and wet patterns change and severe weather occurs with more frequency. The field of quantitative genetics requires climate modeling integration to estimate how genotypes will perform under future climate predictions (Munaweera et al., 2022).

The natural Eucalyptus populations face direct danger from shifts in their native ranges caused by climate change. Current conservation efforts need to use habitat forecast models to protect population areas situated at their range edges. The successful management of climate change impacts will heavily depend on assisted migration practices and ex situ conservation strategies.

Suggestions for Future Research

To address these challenges and advance the field of Eucalyptus quantitative genetics, future research should focus on the following areas:

- **Integration of Genomics and Phenotyping:** Optimal trait measurement precision and genetic selection become achievable by integrating high-resolution genomic data with drone-based image analysis techniques and near-infrared spectral assessment. Large-scale investigations of complex traits will become possible through phenomics platform implementations.
- **Development of Predictive Models:** Machine learning combined with artificial intelligence technologies can help create predictive models for selecting traits in multi-variable frameworks alongside the analysis of gene-environmental elements which results in better breeding across the population. The predictive

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

models should utilize diverse dynamic environmental variables to represent changes in climate patterns.

- **Expansion of Genomic Resources:** Sequencing different Eucalyptus species and populations needs to gain priority because it helps establish a broad pangenome base. Such containment strategies will allow researchers to find rare genetic elements and structural rearrangements which have significance for key traits.
- **Climate-Resilient Breeding:** The selection of breeding traits must focus on characteristics that build climate resiliency among plantings by featuring both drought tolerance and pest resistance abilities. Using adaptive genetic traits from natural populations in breeding programs will produce commercial varieties that show stronger resistance to harsh conditions.

Quantitative genetics drives Eucalyptus breeding while ensuring genetic diversity conservation and developing high-performing new varieties. Research investment in current obstacles combined with emerging technological solutions will expand Eucalyptus capabilities as a leading sustainable forestry and ecological conservation practice in global change scenarios.

CONCLUSION

Quantitative genetics combined with advanced genomic technology now reshapes Eucalyptus breeding while setting the path for tackling both industrial needs and ecological objectives. Modern Eucalyptus breeding programs thrive because they utilize high-throughput sequencing and genomic selection (GS) with marker-assisted selection (MAS) to select traits including growth rate and wood quality together with disease resistance and climate adaptability. The complete genome assembly of Eucalyptus grandis along with EUChip60K SNP arrays combined with predictive models like GBLUP have sped up genetic advancement and cut breeding cycle durations. Natural populations retain their adaptive ability to environmental transformations through enhanced conservation approaches that use knowledge of genetic variability and adaptive gene variants alongside G×E relationships. Genetic improvement efforts face ongoing obstacles which include optimizing multiple traits alongside maintaining robust genomic tools while adapting to climate change scenarios. The upcoming wave of pangenomics technology combined with single-cell sequencing and machine learning methods and advanced phenotyping solutions promises to resolve these barriers. Future genetic research must unify genetic analysis techniques with phenotype evaluation methods to build predictive examination models for complex traits and create abundant genomic assets for improved breeding precision and sustainable long-term objectives. Quantitative genetics establishes itself as an essential force to advance Eucalyptus species through its integration of novel scientific methods and real-world uses. Eucalyptus research maintains industrial compatibility with ecological needs by appropriately linking breeding targets to conservation considerations thus creating sustainable forestry management systems that adapt to environmental change.

Online ISSN 3007-3197
Print ISSN 3007-3189

REFERENCES

Aguirre, N. C., Villalba, P. V., García, M. N., Filippi, C. V., Rivas, J. G., Martínez, M. C., ... & Marcucci Poltri, S. N. (2024). Comparison of ddRADseq and EUChip60K SNP genotyping systems for population genetics and genomic selection in Eucalyptus dunnii (Maiden). Frontiers in Genetics, 15, 1361418.

Ahmar, S., Ballesta, P., Ali, M., & Mora-Poblete, F. (2021). Achievements and challenges of genomics-assisted breeding in forest trees: From marker-assisted selection to genome editing. International Journal of Molecular Sciences, 22(19), 10583.

Alemu, A., Åstrand, J., Montesinos-Lopez, O. A., y Sanchez, J. I., Fernandez-Gonzalez, J., Tadesse, W., ... & Chawade, A. (2024). Genomic selection in plant breeding: Key factors shaping two decades of progress. Molecular Plant.

Amaral, J. S. C. (2021). Decifrando as Respostas de Defesa de Pinus Contra a Infeção por Fusarium Circinatum: Da Fisiologia às Ómicas (Doctoral dissertation, Universidade de Aveiro (Portugal)).

Amrutha, S., Parveen, A. B. M., Muthupandi, M., Vishnu, K., Bisht, S. S., Sivakumar, V., & Ghosh Dasgupta, M. (2021). Characterization of Eucalyptus camaldulensis clones with contrasting response to short-term water stress response. Acta Physiologiae Plantarum, 43, 1-13.

Anilkumar, C., Sunitha, N. C., Harikrishna, Devate, N. B., & Ramesh, S. (2022). Advances in integrated genomic selection for rapid genetic gain in crop improvement: a review. Planta, 256(5), 87.

Baldi, P., & La Porta, N. (2022). Toward the genetic improvement of drought tolerance in conifers: An integrated approach. Forests, 13(12), 2016.

Bartholomé, J., Brachi, B., Marçais, B., Mougou-Hamdane, A., Bodénès, C., Plomion, C., ... & Desprez-Loustau, M. L. (2020). The genetics of exapted resistance to two exotic pathogens in pedunculate oak. New Phytologist, 226(4), 1088-1103.

Begna, T. (2021). Combining ability and heterosis in plant improvement. Open Journal of Plant Science, 6(1), 108-117.

Boursiac, Y., Protto, V., Rishmawi, L., & Maurel, C. (2022). Experimental and conceptual approaches to root water transport. Plant and Soil, 478(1), 349-370.

Bradbury, D., Binks, R. M., & Byrne, M. (2021). Genomic data inform conservation of rare tree species: clonality, diversity and hybridity in Eucalyptus series in a global biodiversity hotspot. Biodiversity and Conservation, 30(3), 619-641.

Bragg, J. G., Yap, J. Y. S., Wilson, T., Lee, E., & Rossetto, M. (2021). Conserving the genetic diversity of condemned populations: Optimizing collections and translocation. Evolutionary Applications, 14(5), 1225-1238.

Butler, J. B., Harrison, P. A., Vaillancourt, R. E., Steane, D. A., Tibbits, J. F., & Potts, B. M. (2022). Climate adaptation, drought susceptibility, and genomic-informed predictions of future climate refugia for the Australian forest tree Eucalyptus globulus. Forests, 13(4), 575.

Calazans, C. C., Souza, J. L., Nunes, V. V., & Silva-Mann, R. (2023). Innovating Eucalyptus resistance: A meta-analysis of gene associations for combatting Leptocybe invasa.

Callister, A. N., Costa-Neto, G., Bradshaw, B. P., Elms, S., Crossa, J., & Brawner, J.

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

T. (2024). Enviromic prediction enables the characterization and mapping of Eucalyptus globulus Labill breeding zones. Tree Genetics & Genomes, 20(1), 3.

Chaudhary, D., Pal, N., Arora, A., Prashant, B. D., & Venadan, S. (2024). Plant Functional Traits in Crop Breeding: Advancement and Challenges. In Plant Functional Traits for Improving Productivity (pp. 169-202). Singapore: Springer Nature Singapore.

Chaves, S. F., Dias, L. A., Alves, R. S., Ferreira, F. M., Araújo, M. S., Resende, M. D., ... & Dias, K. O. G. (2024). Realized genetic gain with reciprocal recurrent selection in a Eucalyptus breeding program. Tree Genetics & Genomes, 20(6), 47.

Climent, J., Alía, R., Karkkainen, K., Bastien, C., Benito-Garzon, M., Bouffier, L., ... & Cavers, S. (2024). Trade-offs and trait integration in tree phenotypes: consequences for the sustainable use of genetic resources. Current Forestry Reports, 10(3), 196-222.

da Silva, P. H. M., Rocha, G. N. D., Araujo, M., Pires Nunes, A. C., & de Paula, R. C. (2024). Thinning Strategies to Optimize Genetic Gain and Population Size in Eucalyptus pellita Breeding. Tree Genetics & Genomes, 20(6), 43.

Dasgupta, M. G., Bari, M. P. A., Shanmugavel, S., Dharanishanthi, V., Muthupandi, M., Kumar, N., ... & Rajasugunasekar, D. (2021). Targeted re-sequencing and genome-wide association analysis for wood property traits in breeding population of Eucalyptus tereticornis× E. grandis. Genomics, 113(6), 4276-4292.

De Mori, G., & Cipriani, G. (2023). Marker-assisted selection in breeding for fruit trait improvement: A review. International Journal of Molecular Sciences, 24(10), 8984.

Dhakal Poudel, P., Cowan, M., Shaw, L., De Faveri, J., Topp, B., & Alam, M. (2023). Macadamia Breeding for Reduced Plant Vigor: Progress and Prospects for Profitable and Sustainable Orchard Systems. Sustainability, 15(19), 14506.

Du, C., Sun, P., Cheng, X., Zhang, L., Wang, L., & Hu, J. (2022). QTL mapping of drought-related traits in the hybrids of Populus deltoides 'Danhong'× Populus simonii 'Tongliao1'. BMC Plant Biology, 22(1), 238.

Duarte, D., Jurcic, E. J., Dutour, J., Villalba, P. V., Centurión, C., Grattapaglia, D., & Cappa, E. P. (2024). Genomic selection in forest trees comes to life: unraveling its potential in an advanced four-generation Eucalyptus grandis population. Frontiers in Plant Science, 15, 1462285.

Duncanson, L., Armston, J., Disney, M., Avitabile, V., Barbier, N., Calders, K., ... & Margolis, H. (2021). Aboveground woody biomass product validation good practices protocol.

Espinosa, E., Bautista, R., Larrosa, R., & Plata, O. (2024). Advancements in long-read genome sequencing technologies and algorithms. Genomics, 110842.

Fahey, M. (2023). Retracing the dispersal of rainforest food trees by east Australian Indigenous Peoples with genomic data (Doctoral dissertation, Macquarie University).

Feng, J., Dan, X., Cui, Y., Gong, Y., Peng, M., Sang, Y., ... & Wang, J. (2024).

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

Integrating evolutionary genomics of forest trees to inform future tree breeding amidst rapid climate change. Plant Communications.

Ferguson, S., Bar-Ness, Y. D., Borevitz, J., & Jones, A. (2024). A telomere-to-telomere Eucalyptus regnans genome: unveiling haplotype variance in structure and genes within one of the world's tallest trees. BMC genomics, 25(1), 913.

Fradgley, N., Gardner, K. A., Bentley, A. R., Howell, P., Mackay, I. J., Scott, M. F., ... & Cockram, J. (2023). Multi-trait ensemble genomic prediction and simulations of recurrent selection highlight importance of complex trait genetic architecture for long-term genetic gains in wheat. in silico Plants, 5(1), diad002.

Gautam, K., Thakur, S., & Kumar, R. V. (2024). Molecular Basis of Forest Genetic Resource Improvement: Progress and Prospects. Biotechnological Approaches for Sustaining Forest Trees and Their Products, 151-180.

Getachew, G., Ayalew, A., Kassahun, H., Tesfaye, A., Alene, M., Redae, T., & Siyum, N. (2024). Evaluate the adaptability of Eucalyptus species in the highlands of the Amhara Region. Heliyon.

González, M. (2023). Estimation of genetic parameters for productive traits in Eucalyptus tereticornis and implications for breeding. BOSQUE, 44(3), 493-502.

Grattapaglia, D. (2022). Twelve years into genomic selection in forest trees: climbing the slope of enlightenment of marker assisted tree breeding. Forests, 13(10), 1554.

Gupta, S., & Shankar, R. (2024). Comprehensive analysis of computational approaches in plant transcription factors binding regions discovery. Heliyon, 10(20).

Hasan, M. M., Thomson, P. C., Raadsma, H. W., & Khatkar, M. S. (2024). A Review and Meta-Analysis of Genotype by Environment Interaction in Commercial Shrimp Breeding. Genes, 15(9), 1222.

Hashmi, Q. A., Ahmad, M., Ahmad, E., Tariq, M., Rafeh, A., & Awais, M. (2023). Exploring drought tolerance in wheat: Insights from biochemical, morphological, and physiological responses. Asian Journal of Research in Crop Science, 8(4), 344-360.

Healey, M. A., Warman, R., Mohammed, C., & Lawson, S. A. (2022). Acacia and eucalypt plantation biosecurity in Southeast Asia–a history, and directions for future research and engagement. Australian Forestry, 85(4), 146-160.

Hua, L., Yu, F., Qiu, Q., He, Q., Su, Y., Liu, X., & Li, J. (2023). Dry-season irrigation further promotes the growth of Eucalyptus urophylla× E. grandis plantations under the conventional fertilization. New Forests, 54(6), 1085-1102.

Hurel, A., de Miguel, M., Dutech, C., Desprez-Loustau, M. L., Plomion, C., Rodríguez-Quilón, I., ... & Budde, K. B. (2021). Genetic basis of growth, spring phenology, and susceptibility to biotic stressors in maritime pine. Evolutionary applications, 14(12), 2750-2772.

Hutapea, F. J., Weston, C. J., Mendham, D., & Volkova, L. (2023). Sustainable management of Eucalyptus pellita plantations: A review. Forest Ecology and Management, 537, 120941.

Ibarra, L., Hodge, G., & Acosta, J. J. (2023). Quantitative genetics of a hybrid population of Eucalyptus nitens× Eucalyptus globulus: estimation of genetic

Online ISSN 3007-3197
Print ISSN 3007-3189

parameters and implications for breeding strategies. Forests, 14(2), 381.

Ikeh, H. (2023). The mechanical properties of wood-doped geopolymer (Master's thesis, Itä-Suomen yliopisto).

Jamil, S., Ahmad, S., Shahzad, R., Umer, N., Kanwal, S., Rehman, H. M., ... & Atif, R. M. (2024). Leveraging Multiomics Insights and Exploiting Wild Relatives' Potential for Drought and Heat Tolerance in Maize. Journal of Agricultural and Food Chemistry, 72(29), 16048-16075.

Jeff-Ego, O. S., Drenth, A., Topp, B., Henderson, J., & Akinsanmi, O. A. (2021). Variability and inheritance in macadamia progenies to Phytophthora cinnamomi and P. multivora the causal agents of root rot and stem canker. Plant and Soil, 466, 449-465.

Jones, C. G. (2024). Development of Microcrystal Electron Diffraction Techniques for the Characterization of Small Molecules and Novel Materials. California Institute of Technology.

Jones, G., Liziniewicz, M., Adamopoulos, S., & Lindeberg, J. (2021). Genetic parameters of stem and wood traits in full-sib silver birch families. Forests, 12(2), 159.

Jordan, R., Prober, S. M., Andrew, R., Freeman, J., Kerr, R., Steane, D., ... & Potts, B. (2023). Population Genomics of Eucalypts.

Kim, S. J. (2024). Population Genomic Study of a Eucalyptus bosistoana Breeding Program (Doctoral dissertation, School of Biological Sciences, University of Canterbury).

Kulak, V., Longboat, S., Brunet, N. D., Shukla, M., & Saxena, P. (2022). In vitro technology in plant conservation: Relevance to biocultural diversity. Plants, 11(4), 503.

Kumar, R., Das, S. P., Choudhury, B. U., Kumar, A., Prakash, N. R., Verma, R., ... & Mishra, V. K. (2024). Advances in genomic tools for plant breeding: harnessing DNA molecular markers, genomic selection, and genome editing. Biological Research, 57(1), 80.

Laidig, F., Feike, T., Hadasch, S., Rentel, D., Klocke, B., Miedaner, T., & Piepho, H. P. (2021). Breeding progress of disease resistance and impact of disease severity under natural infections in winter wheat variety trials. Theoretical and Applied Genetics, 134, 1281-1302.

Lee, S. H., Dahali, R., Nik Hashim, N. H., Kusin, M., Mahmud, S. Z., Kamarudin, N., ... & Lubis, M. A. R. (2023). Eucalyptus Plantation Worldwide, Its Hybridization and Cloning Development. In Eucalyptus: Engineered Wood Products and Other Applications (pp. 1-15). Singapore: Springer Nature Singapore.

Li, C., Zhang, L., Peng, Z., Li, X., Liu, Z., Lu, T., ... & Yang, J. (2024). Genetic relationship analysis and core collection construction of Eucalyptus grandis from Dongmen improved variety base: the largest eucalypt germplasm resource in China. BMC Plant Biology, 24(1), 1240.

Li, Z., Zhao, Y., & Luo, K. (2024). Molecular Mechanisms of Heterosis and Its Applications in Tree Breeding: Progress and Perspectives. International Journal of Molecular Sciences, 25(22), 12344.

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

Liu, G., Xie, Y., Shang, X., & Wu, Z. (2021). Expression patterns and gene analysis of the cellulose synthase gene superfamily in Eucalyptus grandis. Forests, 12(9), 1254.

Lott, R., & Read, S. (2021). Status of Australia's Forest Genetic Resources 2021. ABARES Research Report 21.15, Australian Bureau of Agricultural and Resource Economics and Sciences, Canberra.

Lu, Y., & Fricke, W. (2023). Salt Stress—Regulation of root water uptake in a whole-plant and diurnal context. International Journal of Molecular Sciences, 24(9), 8070.

Lyu, X., Li, J., & Li, S. (2024, November). Approaches to Reach Trustworthy Patient Education: A Narrative Review. In Healthcare (Vol. 12, No. 23, p. 2322). MDPI.

Makouanzi, G., Chaix, G., Nourissier, S., & Vigneron, P. (2018). Genetic variability of growth and wood chemical properties in a clonal population of Eucalyptus urophylla× Eucalyptus grandis in the Congo. Southern Forests: a Journal of Forest Science, 80(2), 151-158.

Malakar, M. (2024). Eucalyptus Spp.: A Wonder Tree. In Advances in Medicinal and Aromatic Plants (pp. vol1-185). Apple Academic Press.

Manzoor, S., Nabi, S. U., Rather, T. R., Gani, G., Mir, Z. A., Wani, A. W., ... & Manzar, N. (2024). Advancing crop disease resistance through genome editing: a promising approach for enhancing agricultural production. Frontiers in Genome Editing, 6, 1399051.

Mapari, A. R., & Mehandi, S. (2024). Enhancing Crop Resilience: Advances and Challenges in Marker-Assisted Selection for Disease Resistance. Journal of Advances in Biology & Biotechnology, 27(7), 569-580.

Mathiazhagan, M., Chidambara, B., Hunashikatti, L. R., & Ravishankar, K. V. (2021). Genomic approaches for improvement of tropical fruits: fruit quality, shelf life and nutrient content. Genes, 12(12), 1881.

McBreen, J., Babar, M. A., Jarquin, D., Khan, N., Harrison, S., DeWitt, N., ... & Van Sanford, D. A. (2025). Enhancing prediction accuracy of grain yield in wheat lines adapted to the southeastern United States through multivariate and multi-environment genomic prediction models incorporating spectral and thermal information. The Plant Genome, 18(1), e20532.

McGowan, M., Wang, J., Dong, H., Liu, X., Jia, Y., Wang, X., ... & Zhang, Z. (2021). Ideas in genomic selection with the potential to transform plant molecular breeding: a review. Plant breeding reviews, 45, 273-319.

Mieres-Castro, D., Maldonado, C., & Mora-Poblete, F. (2024). Enhancing prediction accuracy of foliar essential oil content, growth, and stem quality in Eucalyptus globulus using multi-trait deep learning models. Frontiers in Plant Science, 15, 1451784.

Misztal, I., Aguilar, I., Lourenco, D., Ma, L., Steibel, J. P., & Toro, M. (2021). Emerging issues in genomic selection. Journal of animal science, 99(6), skab092.

Müller, B. S., Neves, L. G., de Almeida Filho, J. E., Resende, M. F., Muñoz, P. R., Dos Santos, P. E., ... & Grattapaglia, D. (2017). Genomic prediction in contrast to a

Online ISSN 3007-3197
Print ISSN 3007-3189

genome-wide association study in explaining heritable variation of complex growth traits in breeding populations of Eucalyptus. BMC genomics, 18, 1-17.

Munaweera, T. I. K., Jayawardana, N. U., Rajaratnam, R., & Dissanayake, N. (2022). Modern plant biotechnology as a strategy in addressing climate change and attaining food security. Agriculture & Food Security, 11(1), 1-28.

Negash, L. (2021). A Selection of African Native Trees: Biology, Uses, Propagation and Restoration Techniques. Addis Ababa, Ethiopia.

O'Connell, A., Bolwell, P., & Schuch, W. (2022). Impact of forest tree biotechnology on the pulp and paper-making processes in the 21st century. In Transgenic plant research (pp. 175-186). Routledge.

Ogunbawo, A. R., Mulim, H. A., Campos, G. S., Schinckel, A. P., & Oliveira, H. R. D. (2024). Tailoring Genomic Selection for Bos taurus indicus: A Comprehensive Review of SNP Arrays and Reference Genomes. Genes, 15(12), 1495.

Olaoye, K. O., & Ojo, M. O. (2022). Non-destructive acoustic assessment of wood quality in trees and logs and the effects of silvicultural treatments: a review. International Wood Products Journal, 13(3), 156-171.

Paludeto, J. G. Z., Grattapaglia, D., Estopa, R. A., & Tambarussi, E. V. (2021). Genomic relationship–based genetic parameters and prospects of genomic selection for growth and wood quality traits in Eucalyptus benthamii. Tree Genetics & Genomes, 17(4), 38.

Persa, R., Ribeiro, P. C. D. O., & Jarquin, D. (2021). The use of high-throughput phenotyping in genomic selection context. Crop Breeding and Applied Biotechnology, 21(spe), e385921S6.

Pfeilsticker, T. R., Jones, R. C., Steane, D. A., Harrison, P. A., Vaillancourt, R. E., & Potts, B. M. (2022). Expansion of the rare Eucalyptus risdonii under climate change through hybridization with a closely related species despite hybrid inferiority. Annals of Botany, 129(1), 1-14.

Pinheiro, A. L. M. (2021). Coexpression network analysis of proteins and metabolites time-course of two contrasting Eucalyptus grandis reponses to Austropuccinia psidii (Doctoral dissertation, Universidade de São Paulo).

Prober, S. M., Potts, B. M., Harrison, P. A., Wiehl, G., Bailey, T. G., Costa e Silva, J., ... & Vaillancourt, R. E. (2022). Leaf economic and hydraulic traits signal disparate climate adaptation patterns in two co-occurring woodland eucalypts. Plants, 11(14), 1846.

Ribeiro, F. P., Gatto, A., Oliveira, A. D. D., Pulrolnik, K., Valadão, M. B. X., Araújo, J. B. C. N., ... & Ferreira, E. A. B. (2023). Carbon Storage in Different Compartments in Eucalyptus Stands and Native Cerrado Vegetation. Plants, 12(14), 2751.

Rocha, L. F., Benatti, T. R., de Siqueira, L., de Souza, I. C. G., Bianchin, I., de Souza, A. J., ... & Tambarussi, E. V. (2022). Quantitative trait loci related to growth and wood quality traits in Eucalyptus grandis W. Hill identified through single- and multi-trait genome-wide association studies. Tree Genetics & Genomes, 18(6), 38.

Rocha-Sepúlveda, M. (2023). Genetic and environmental controls of traits affecting profitability of pulp and solid-wood production from plantation-grown

Online ISSN 3007-3197
Print ISSN 3007-3189

Eucalyptus nitens (Doctoral dissertation, PhD thesis, University of Tasmania, Hobart, Tas., Australia).

Sandhu, K. S., Shiv, A., Kaur, G., Meena, M. R., Raja, A. K., Vengavasi, K., ... & Kumar, S. (2022). Integrated approach in genomic selection to accelerate genetic gain in sugarcane. Plants, 11(16), 2139.

Serpico, D., Lynch, K. E., & Porter, T. M. (2023). New historical and philosophical perspectives on quantitative genetics. Studies in History and Philosophy of Science, 97, 29-33.

Shahrajabian, M. H., Sun, W., & Cheng, Q. (2021). Molecular breeding and the impacts of some important genes families on agronomic traits, a review. Genetic Resources and Crop Evolution, 68(5), 1709-1730.

Sharma, P., Doultani, S., Hadiya, K. K., George, L. B., & Highland, H. N. (2024). Overview of Marker-assisted Selection in Animal Breeding. Journal of Advances in Biology & Biotechnology, 27(5), 303-318.

Sharma, U., Sankhyan, H. P., Kumari, A., Thakur, S., Thakur, L., Mehta, D., ... & Sankhyan, N. (2024). Genomic selection: a revolutionary approach for forest tree improvement in the wake of climate change. Euphytica, 220(1), 9.

Singh, M., Nara, U., Kumar, A., Thapa, S., Jaswal, C., & Singh, H. (2021). Enhancing genetic gains through marker-assisted recurrent selection: from phenotyping to genotyping. Cereal Research Communications, 1-16.

Sinha, D., Maurya, A. K., Abdi, G., Majeed, M., Agarwal, R., Mukherjee, R., ... & Chen, J. T. (2023). Integrated genomic selection for accelerating breeding programs of climate-smart cereals. Genes, 14(7), 1484.

Sofi, P. A., Rehman, K., Gull, M., Kumari, J., Djanaguiraman, M., & Prasad, P. V. V. (2021). Integrating root architecture and physiological approaches for improving drought tolerance in common bean (Phaseolus vulgaris L.). Plant Physiology Reports, 26, 4-22.

Souden, S., Ennajeh, M., Ouledali, S., Massoudi, N., Cochard, H., & Khemira, H. (2020). Water relations, photosynthesis, xylem embolism and accumulation of carbohydrates and cyclitols in two Eucalyptus species (E. camaldulensis and E. torquata) subjected to dehydration–rehydration cycle. Trees, 34, 1439-1452.

Tang, M., Wang, T., & Zhang, X. (2022). A review of SNP heritability estimation methods. Briefings in bioinformatics, 23(3), bbac067.

Thapliyal, G., Bhandari, M. S., Vemanna, R. S., Pandey, S., Meena, R. K., & Barthwal, S. (2023). Engineering traits through CRISPR/cas genome editing in woody species to improve forest diversity and yield. Critical Reviews in Biotechnology, 43(6), 884-903.

Tomé, M., Almeida, M. H., Barreiro, S., Branco, M. R., Deus, E., Pinto, G., ... & Rodríguez-Soalleiro, R. (2021). Opportunities and challenges of Eucalyptus plantations in Europe: the Iberian Peninsula experience. European Journal of Forest Research, 140(3), 489-510.

Trudić, B., Stojnić, S., Avramidou, E. V., Malliarou, E., Ivezić, A., Bojović, M., & Pilipović, A. (2024). Recent global (bio) technological advancements in the service of forest genetic resources protection under climate change. časopis Topola, (213), 17-31.

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

Udoh, L. I., Obaseojei, W. P., & Uzoebo, C. (2021). Single nucleotide polymorphisms: a modern tool to screen plants for desirable traits. In Plant breeding-current and future views. IntechOpen.

Valenzuela, C. E., Ballesta, P., Ahmar, S., Fiaz, S., Heidari, P., Maldonado, C., & Mora-Poblete, F. (2021). Haplotype-and SNP-based GWAS for growth and wood quality traits in Eucalyptus cladocalyx trees under arid conditions. Plants, 10(1), 148.

Varghese, M., Kamalakannan, R., & Suraj, P. G. (2024). Eucalyptus: Taxonomy, Geographic Distribution, Domestication, Breeding, Ecology and Economic Importance. In Economically Important Trees: Origin, Evolution, Genetic Diversity and Ecology (pp. 81-128). Springer, Singapore.

Wang, J., Hou, Y., Wang, Y., & Zhao, H. (2021). Integrative lncRNA landscape reveals lncRNA-coding gene networks in the secondary cell wall biosynthesis pathway of moso bamboo (Phyllostachys edulis). BMC genomics, 22, 1-13.

Wang, M., Li, R., & Zhao, Q. (2023). Multi-omics techniques in genetic studies and breeding of forest plants. Forests, 14(6), 1196.

Wang, W. (2024). Genome Editing Improvement Study of Eucalyptus Wood Quality Traits. Molecular Plant Breeding, 15.

Wang, Y., & Zhang, L. (2024). Gene-driven future: breakthroughs and applications of marker-assisted selection in tree breeding. Molecular Plant Breeding, 15.

Wang, Y., Zafar, N., Ali, Q., Manghwar, H., Wang, G., Yu, L., ... & Jin, S. (2022). CRISPR/Cas genome editing technologies for plant improvement against biotic and abiotic stresses: advances, limitations, and future perspectives. Cells, 11(23), 3928.

Xiao, Q., Bai, X., Zhang, C., & He, Y. (2022). Advanced high-throughput plant phenotyping techniques for genome-wide association studies: A review. Journal of advanced research, 35, 215-230.

Xing, Y., & Wang, X. (2024). Precision agriculture and water conservation strategies for sustainable crop production in arid regions. Plants, 13(22), 3184.

Younessi-Hamzekhanlu, M., & Gailing, O. (2022). Genome-wide SNP markers accelerate perennial forest tree breeding rate for disease resistance through marker-assisted and genome-wide selection. International Journal of Molecular Sciences, 23(20), 12315.

Yunus, M. H., Firdaus, A., Khan, Z., & Ansari, M. Y. K. (2025). 7 Genomics-Assisted Breeding (GAB) for Trait Improvement: Unveiling Genomic Strategies for Accelerated Crop Enhancement. Plant Breeding Technology: Future Trends and Challenges, 138.

Zampieri, E., Pesenti, M., Nocito, F. F., Sacchi, G. A., & Valè, G. (2023). Rice Responses to Water Limiting Conditions: Improving Stress Management by Exploiting Genetics and Physiological Processes. Agriculture, 13(2), 464.

Zhao, Y., Tian, Y., Sun, Y., & Li, Y. (2022). The Development of Forest Genetic Breeding and the Application of Genome Selection and CRISPR/Cas9 in Forest Breeding. Forests, 13(12), 2116.

Zhu, X., Weng, Q., Bush, D., Zhou, C., Zhao, H., Wang, P., & Li, F. (2023). High-density genetic linkage mapping reveals low stability of QTLs across

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

environments for economic traits in Eucalyptus. Frontiers in Plant Science, 13, 1099705.

al.

The Margalla Hill Limestone and Chorgali Formation plays a vital role in the interpretation of carbonate sedimentation in the area, and the manner in which shallow marine environment dominated during the Early Eocene (DiPietro & Pogue, 2004). These formations consist mainly of carbonates and are interleaved with shales and marls, which provide evidence of varying environmental conditions during the Eocene (Jain, 1981). The faunal assemblages of large benthic foraminifera such as, Nummulites and Assilina are a rich feature of the Margalla Hill Limestone that are used as important biostratigraphic markers for determining age and palaeogeographic reconstruction (Shah, 1977; Munir et al., 2006). The overlying Chorgali Formation over the Margalla Hill Limestone also contributes towards the marine ecosystem of the area and the faunal assemblies of its time.

The deposition comparing with the sedimentary environment and the sedimentary process, which related to the forming of these carbonates, is an important clue to the restoring of paleogeography. The microfacies analysis of Margalla Hill Limestone and Chorgali Formation provides a clue about the shallow marine environments which existed during the early Eocene (Singh, 1991). Microfacies approach makes a possible to recognize proper assemblages and polution a type described above as mo-compatible with dynamic occuring syngenetic environmental phenomena during sedimentation of these carbonates. Interpretations for these microfacies types will be used to demarcate the inner ramp, middle ramp, and lagoon depositional environments and to infer energy regime and paleogeography during the time.

The main objective is to interpret the lithofacies of the Margall Hill Limestone and Chorgali Formation & to determine the depositional environments on the basis of Microfacies types identified. This investigation will improve our depositional model for the Early Eocene carbonates of the Balakot area, thus, enhancing our knowledge regarding the carbonate sedimentation in the shallow marine environment. The research in result will offer significant information on the regional tectonic history and paleogeographic evolution of the area during the Himalayan orogeny stage-Tectonic and environmental – nonlinear changes.

Research Objectives

1. Identification and classification of microfacies types in the Margalla Hill Limestone and Chorgali Formation, including foraminiferal bioclastic wackestones, Assilina-Nummulites bioclastic wackestones, and bioclastic mudstones.
2. Interpretation of the depositional environments of these carbonates, specifically inner

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

ramp, middle ramp, and lagoonal settings, based on the identified microfacies.

3. Discussion of sedimentary processes that shaped the Early Eocene carbonates in the context of tectonic and paleogeographic changes in the region.

1. Geological Setting

Regional Geology

The tectonically active Sub-Himalayan realm, formed due to the constructive plate boundary, known as the Himalayan orogenic belt (Molnar and Tapponier, 1975) envelops the northern Pakistan's Balakot area in it. The geological setting of this area is attributed to collision between the Indian and Eurasian plate, which started from the Late Cretaceous, and currently is still the actuation of fold-thrust belts, fault systems, and sedimentary basins (Yin, 2006). The occurrence of the MBT and other major faulting and folding on the northwestern limb of the Hazara-Kashmir Syntaxis is a significant feature in terms of the uplift and facial exposure of rocks and separation of older and younger strata such as Margalla hill Limestone and Chorgali Formation.




Figure 1: Tectonic Map of Northern Pakistan

This map illustrates the major structural boundaries and tectonic divisions in the region. It would complement your discussion of the tectonic setting, particularly the reference to the Balakot region's position in the Sub-Himalayan zone and its relationship with the MBT

Stratigraphy of the Study Area

The Balakot region's stratigraphy is dominated by Cenozoic deposits, particularly the Eocene units, which include the Margalla Hill Limestone and Chorgali Formation. These formations provide insight into Early Eocene depositional environments and tectonic history:

1. *Abbottabad Formation (Cambrian):* Marine dolomites, sandstones, and shales.
2. *Hangu Formation (Lower Paleocene):* Sandstones and carbonaceous shales, indicating

Online ISSN 3007-3197
Print ISSN 3007-3189

shallow marine environments.

3. *Lockhart Formation (Late Paleocene):* Limestones and calcareous sandstones, containing significant fossils.
4. *Patala Formation (Early Eocene*)**:** Shales and limestones from a shallow marine setting.
5. *Margalla Hill Limestone (Middle Eocene):* Nodular limestone with abundant larger benthic foraminifera, crucial for biostratigraphy (Shah, 1977).
6. *Chorgali Formation (Upper Eocene):* Nodular limestone with foraminiferal marls and shales, offering insights into paleogeographic history.

These units reflect dynamic depositional environments and sea-level changes, with the Margalla Hill Limestone and Chorgali Formation being key to understanding Early Eocene transgressive-regressive cycles and paleoclimatic conditions.