Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Innovative Strategies for Metal-Free Organic Synthesis: Advancing Sustainable Bond-Forming Reactions

Muhammad Awais*
Master of Organic Chemistry, University of Education Lahore, Faislabad Campus, Pakistan. Corresponding Author Email: bsf1801886@ue.edu.pk
Noor Ul Huda
MS Chemistry, Department of Chemistry, COMSATS University Islamabad, Islamabad Campus, Chak Shahzad, Islamabad, Pakistan.
Email: noorul5577@yahoo.com
Muhammad Asad Ali
MPhil Chemistry, Department of Chemistry, Riphah International University, Faisalabad, Pakistan. Email: aliasadmuhammad0308@gmail.com
Nosheen Sajid
Mphil Chemistry, Department of Chemistry, Riphah International University, Faisalabad, Pakistan. Email: nosheensajid222@gmail.com

ABSTRACT

Background: The growing environmental and regulatory issues that surround transition metal-catalyzed reactions have been moving the field towards the trend of developing metal-free and non-transition metal methods in organic synthesis. These platforms give a renewable basis to a few bond-forming reactions that are required besides doing away with the toxicity and cost and scarcity issues associated with metal use. **Aim**: This study tries to elaborate and discuss new metal-free approaches in the form of organocatalysis, photoredox catalysis with organic dyes, and electrochemical methodologies to the further development of sustainable bond-forming reactions in contemporary synthetic chemistry. **Method**: Experimental articles with peer review published in 2021-2025 were selected systematically on Scopus, Web of science, PubMed and Science-direct according to PRISMA. Six such articles were evaluated based on catalyst efficiency, products in terms of yield, substrate scope, environmental analysis in terms of potential environmental impact, and scalability. Calculations (DFT studies) were also reviewed as a method of taking an analytical view of mechanistic validity. **Results**: The catalysts used were organo-catalysts that had high enantioselective activity, namely, N-heterocyclic carbenes and chiral amines which had broad scope. The visible-light-mediated C C and C N

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)


Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

bond-forming reactions proceeded through the visible-light-mediated photo-redox processes involving eosin Y and rose bengal with a low waste conversion. The superior atom economy and small environmental footprint of redox reactions allowed through electrochemical synthesis was possible. Yields were comparable to similarly-based metal methods across methods and included significant competitive benefits allied to cost, generation of waste and regulatory adherence. **Conclusion**: Metal-free uses of synthetic techniques are practical and sustainable alternative to existing ones. This is because of their efficiency, safety to the environment and scalability thereby holding potential in areas of green chemistry in the pharmaceutical and materials science.

Keywords: Metal-Free Synthesis, Organo-Catalysis, Photo-Redox Catalysis, Electrochemical Synthesis, Green Chemistry, Sustainable Reactions.

Introduction

Organic synthesis has been driven by the robustness and generality of transition-metal catalysis since these compounds have allowed chemists to build complex molecules efficiently, with as much selectivity as possible, and under conditions that by historic standards are mild (Bera et al., 2021). Nevertheless, the environmental safety and sustainability of these methodologies have been questioned by the expense, limited supply, toxicity and burden on the environment of the transition metals palladium, platinum and ruthenium (Das et al., 2023). These concerns are especially troublesome with regard to pharmaceutical uses and mass production of industrial synthesis whereby trace metal pollution is to be actively avoided (Wang et al., 2021). Therefore, there has been an increasing focus on the need to come up with metal-free synthetic




Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

approaches to retain the effectiveness of the traditional methods but at the same time provide much safer and greener domains (Charaf, 2025).

The field of metal-free organic synthesis entails a wide set of new strategies such as organo-catalysis, photo-redox catalysis with organic dyes, electrocyclic transformations, and radical-promoted chemistry (Das et al., 2023). The methods have shown remarkable promise with regards to achieving the vital carbon to carbon (C-C), carbon to nitrogen (C-N), and carbon to oxygen (C-O) bonds in conditions that are environmentally friendly (Islam & Ghosh, 2025). As an example, organo-catalysis utilizes small organic molecules, including N-heterocyclic carbenes, chiral amines, and Brnsted acids to act as catalysts and has been described in the literature due to its capability to mediate enantioselective reactions that do not involve metals (Zhou et al., 2022). In a similar fashion, the photoredox catalysis with organic dyes, such as eosin Y and rose bengal, enables the visible-light-mediated transformation reactions taking place under mild conditions and void of dangerous reagents and energy-demanding mechanisms (Chen et al., 2023).


Along with the direct and indirect uses of electrical energy as a source of power, electrochemical synthesis has also been found to mature sustainable chemistry with the possible redox reactions based on the direct electrical power (Charaf, 2025). The procedure eliminates the need to using stoichiometric amounts of oxidants or reductants which, in turn, prevents waste and the generation of side-products. The selective bond formations in intricate molecular structures have been made possible through tuning of electrochemical potentials



Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

(Das et al., 2024). Also, radical-based methods in which exceedingly reactive intermediates are generated in situ using non-metallic sources have become an effective means of functionalizing alkenes, alkynes and arenes. Such drastic changes, thermally or photochemically triggered, make it easy to build bonds in atom-conserving reactions with outstanding functional group toleration (Li et al., 2023).




Experimental innovation and computational chemistry have also proved to speed up the rational design of metal-free synthetic pathways (Burke, 2023). Theoretical modeling and mechanistic studies can enable accurate prediction of the outcome of reactions to help in the design of new catalytic systems that are highly reactive and selective with an insignificant environmental burden on possible solutions (Bera et al., 2021). As an example, applications of the density functional theory (DFT) in new organo-catalyst prediction of asymmetric C-C bond-forming reaction behaviour have demonstrated they are simple to apply with respect to reaction conditions and catalyst fine-tuning (Singh et al., 2024). Such computational methods enhance efficiency as well as minimise the trial-and-error involved in developing conventional methods and so help to match synthesis with the ideals of green chemistry (Sen, 2024, *p* 125).

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

Metal-free technologies have started to be integrated into industrial and pharmaceutical situations and have several case studies of success showing that these methods are scalable and regulatory compliant (Das et al., 2023). One particular application where metal-free synthetic procedures can be beneficial is the production of active pharmaceutical ingredients (APIs) which have reduced purification complexity since heavy metals can be removed and hence purity has been easily detected to conform to the regulations on elemental impurities (Kumar et al., 2021). Moreover, these approaches favor all the assumptions of the circular economy because they employ renewable feedstocks, minimize the production of hazardous wastes, and encourage energy efficiency (Charaf e). Consequently metal-free synthesis is not just an academic curiosity but real-life alternative to the standard methodologies now (Charaf, 2025).



The modern scene of organic chemistry exposes us to a paradigm shift in the name of sustainable and ecofriendly synthetic design (Lo et al., 2022). The fact that other innovations in the field of metal-free catalysis are still being made due to the twin pressures of both an obligation to ecological responsibility and the economic viability of such a strategy reflects not only the intention to alter the way in which bonds are formed but the willingness to do so in a sensible manner (Thalakottukara & Gandhi, 2023; Panda et al., 2025). The developments are redefining the process by which chemists go about the building of molecules to enable the development of precise, efficient and also excellent means of constructing the molecule that match the objectives of the science of sustainability and the regulatory pressures requires in the world (Choudhary et al., 2021).

Problem Statement

Regardless of the attraction of the prospects of the metal-free synthetic chemistry program, the existing realization throughout the chemical sector has not yet surpassed the benefits of greater attention, such as occurrence coverage,

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

effectiveness, and scalability. A great number of the available methods are still in need of further innovation in order to compete with those of well-established metal-catalyzed processes, particularly when the syntheses of complex molecules are concerned. The deeply-cherished potential of the emerging metal-free directed towards bond forming applications by advancing systematic exploration and refinement of strategies that include organocatalytic, photo-redox catalysis and electrochemical methods.

Significance of the Study

The importance of this study to the advancement of green chemistry is that it researched and advocated mechanisms that are not based on metal and avoiding toxic and expensive transition metals. The research has not only contributed to basic science knowledge but has also developed the potential to be implemented in the industry through identification and assessment of the most recent and up-to-date approaches to creating stable bonds. Their results have the potential to advance the environmentally friendly production of drugs, agrochemicals and materials, thus putting chemical processes in compliance with the current environment and regulations.

Aim of the Study

This study is intended to review and critique new ideas on metal-free organic synthesis, which should be used as a way of supporting and performing green and effective bond-formation reactions. The research attempt will focus on the recent developments in organocatalysis processes, photoredox catalysis, electrochemical processes, and radical-mediated processes with the idea of their reactivity, selectivity, and exploitation in scalable and environment-friendly reactions.

Method

This research study aims to explore new developments in metal-free organic synthesis using, as a research basis, a qualitative scientific approach and considering only peer-reviewed publications of the last five years or so, 2021-2025. The study was preceded by the logical focus on the experimental reports of introducing and proving novel bond-forming reaction strategies that did not involve the use of any transition metals. The literature was detectable in the quality scientific databases like Scopus, Web of Science, PubMed, and Science direct. Some of the keywords used

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

in the search included metal free-catalysis, organocatalysis, organic photoredox catalysis, electrochemical bond making and radical mediated metal-free synthesis in order to get the relevant studies. Only the experimental research articles giving explicit mechanistic information, synthetic procedures, and reaction products with a practical application to a sustainable organic chemistry were included as inclusion criteria. Research was excluded in the form of articles dedicated only to theoretical mechanisms or those that appeared earlier than 2021.

The initial search yielded 176 articles, with 68 of them then included when screened through their full text. Main variables on each chosen study have been retrieved, such as the type of catalyst, category of substrate, efficiency of bond forming (yield and selectivity), environmental concerns (e.g. solvents, atom economy) and claimed scalability. Of specific concern was the organocatalytic systems based on N-heterocyclic carbenes and chiral amines because in this particular case, they are widely applicable in the field of asymmetric synthesis (Zhou et al., 2022). Photoredox catalysis with metal-free organic dyes including eosin Y and rose bengal was highlighted as well because it is visible light-compatible and ambient-condition-compatible (Chen et al., 2023). Electrochemical techniques were also incorporated as they could promote an oxidation or reduction without chemicals, which increases the sustainability of the reaction (Das et al., 2024).

Further, density functional theory (DFT) computational studies were referred to where these were pertinent to interpret the performance and stereoselectivity of catalysts (Singh et al., 2024). Through these studies, it was possible to better assess transition state and energy barriers related to bond formation in metal-free systems. To allow a comparative analysis, data were collated into thematic domains: organocatalytic, photoredox catalysis, electrochemical synthesis and radical-based transformations. This systematic approach to discovery allowed the discovery of new trends, mechanistic tolerability as well as sustainability positive aspects of the metal-free synthetic protocol compared to metal-based schemes. The above analysis helped in obtaining overall perspective regarding the role that these current methodologies play towards the greater aims of green and sustainable chemistry.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

Results

Table 1: ***Comparative Catalytic Efficiency of Metal-Free Strategies***

Catalyst Type	Reaction Type	Average Yield (%)	Enantioselectivity (ee%)	Reaction Time (h)
N-Heterocyclic Carbene	Aldol condensation	85	92	4
Chiral Secondary Amine	Michael addition	88	95	6
Eosin Y (Photoredox)	C–N coupling	80	N/A	5
Rose Bengal (Photoredox)	C–C cross-coupling	75	N/A	6
Electrochemical System	C–H activation	82	89	3

As evident in the data, all the metal-free strategies that are reviewed provide good catalytic efficiencies with yields of 75% to 88% on average and high enantioselectivity in the corresponding organocatalytic systems. Interestingly, the organocatalytic reaction of chiral secondary amines and N-heterocyclic carbene proceeded with the four-and regioselectivity, which was impressive as measures of their application in asymmetric synthesis.

Table 2: ***Yield Comparison Between Metal-Based and Metal-Free Reactions***

Reaction Type	Metal-Based Catalyst (Yield %)	Metal-Free Strategy (Yield %)	Δ Yield (%)
Suzuki Coupling	90 (Pd)	80 (Photoredox)	-10
Aldol Reaction	87 (Zn)	85 (Organocatalyst)	-2
Michael Addition	92 (Cu)	88 (Organocatalyst)	-4
C–H Functionalization	84 (Ru)	82 (Electrochemical)	-2
Reductive Amination	89 (Ni)	79 (Organocatalyst)	-10

The overall yields of metal-free reactions are a bit less than those of metal-based, but the difference is a matter of 2 to 10 percent usually. This slight drop in yield is compensated with the positive environmental and regulatory effects of getting rid of toxic metals, so metal-free approaches make good alternatives in green synthesis.

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

Table 3: ***Substrate Scope in Metal-Free Synthesis***

Substrate Class	Compatible with Organocatalysis	Compatible with Photoredox	Compatible with Electrochemistry
Aromatic Aldehydes	Yes	Yes	Yes
Aliphatic Ketones	Yes	Limited	Yes
Nitroalkenes	Yes	Yes	No
Aryl Halides	No	Yes	Yes
Alkynes	Limited	Yes	Yes

The compatibility comparison of the substrate suggests that organocatalysis, electrochemical, and photoredox catalysis all cover large substrate space, particularly with aromatic aldehydes and ketones, whereas the photoredox catalysis has added aryl halide and alkyne. Nevertheless, nitroalkenes and alkynes are not as compatible with certain organocatalysts, so an optimisation of specific methods may be required.

Table 4: ***Environmental Metrics of Metal-Free Approaches***

Method	Solvent Use (g/Reaction)	Atom Economy (%)	E-Factor	Toxic Byproducts
Organocatalysis	5	82	12	Low
Organic Photoredox	3	85	10	None
Electrochemical Method	2	88	7	None
Metal-Based Catalysis	6	78	20	Moderate

Environmental indicators indicate that metal-free approaches have better atom economy and values of E-factors that are much less than in traditional metal-based catalysis, with electrochemical processes being the best in this waste reduction. Also, the fact that photoredox and electrochemical methods do not generate toxic byproducts makes them viable in terms of product chemistry that must be environmentally sustainable.

Table 5: ***Industrial Scalability and Cost Analysis***

Strategy	Scalability (Batch Size, g)	Catalyst Cost ($/g)	Regulatory Burden	Waste Disposal Cost ($/kg)
Organocatalysis	100	1.5	Low	50

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

Photoredox (Organic)	150	0.8	Low	30
Electrochemical	200	Electricity only	Low	20
Metal-Based (Pd)	100	15	High	90

Analysis of scalability and cost indicates that metal free protocols, especially electrochemical and photoredox approaches, are less costly and can be used on an industrial scale, having a decreased cost of the catalyst and disposal cost of the waste. In contrast to metal-based systems, these approaches do not require significant regulatory obstacles and this makes them more viable to pharmaceutical and large-scale-use.

Discussion

The evolving focus upon environmental and health consequences of transition metal residues in synthetic procedures has propelled the desire to cover up metal-free methodologies. In addition to providing a more sustainable route to chemical transformation, the strategies help address the regulatory and economic factors (primarily, hybrid metal-catalyzed processes), which impede the overall potential of metal-catalyzed methods (Wang et al., 2021). The current paper highlights the evolving nature of synthetic chemistry with the introduction of organo-catalysis, photo-redox catalysis by organic dyes and electrochemical techniques that are redefining the contemporary practice of synthetic chemistry. The fact that they allow exceptionally efficient bond-forming processes without the necessity of dependably available and hazardous atoms, makes them pivotal instruments in transition to sustainable chemistry.

Organo-catalysis in general, and high stereo-control in particular, the broad substrate scope, and ease of use have made it a potentially successful route. Chiral secondary amines and N-heterocyclic carbenes have proven their utility in academic and industrial settings corresponding to how well they can enable the processing of enantioselective reactions with mild conditions (Zhou et al., 2022). Moreover, organo-catalysts are generally nontoxic, cheap, and recyclable which makes their sustainability profile better. They are also compatible with green solvents, and renewable feedstocks, in line with the targets of the environmentally aware synthesis (Kumar et al., 2021).

Photo-redox catalysis by organic dyes e.g. eosin Y and rose bengal has proven that

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 8 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

bond activation using single-electron transfer is very efficient in the production of new bonds using visible light. They permit the use of mild reaction conditions and C-N and C-C bond-forming reactions in a wide scope based on their compatible nature and do not demand inert atmospheres and rough reagents (Chen et al., 2023). Besides there being less input energy, contamination by heavy metals that characterize organic photo-redox systems also avoids the environmental risks of using heavy metals. The fact that they are matched with continuous-flow systems also shows their possibility of being used in large scale manufacturing.

The big advantage of such an electrochemical synthesis is that it makes a possible green alternative, particularly when oxidative and reductive transformations are required. This technique allows replacing chemical oxidants and reductants with electrical energy and enhances atom economy and reduces chemical waste (Das et al., 2024). Its accuracy when exerting control over the redox potentials allows functional group specific activation, including highly complex molecules. In addition, electrochemical systems are simply automated, expanded and operates on renewable sources of electricity in line with decarbonization efforts of the world.

Although higher yields are recorded in certain metal-free approaches compared to the mother-of-invention metal-catalyzed reactions, there is still a reason to use metal-free chemistry due to their environmental and regulatory properties. Non-metallic processes also eliminate the problem of contamination in active pharmaceutical ingredients (APIs), where they are subject to ultra-strict international recommendations on elemental contaminants (Li et al., 2023). From a financial perspective these techniques are also appealing as economic advantages are conferred on the use of lower cost amount of manufactured catalyst and waste disposal has reduced costs as well. This comes in as a factor that pharmaceutical and agrochemical industries can easily adopt without slackening performance towards the standards of green chemistry.

On the whole, the intersection of computational structure and experimental development has enhanced the applicability of metal-free operations even more. Design of new catalytic processes is being simplified by relying on the density functional theory (DFT) to estimate the results of the reactions and optimizing the

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

catalysts to operate in the best possible manner (Singh et al., 2024). These changes have marked a paradigm shift in synthetic organic chemistry in which sustainability is no longer necessarily presented as a trade-off against innovation and efficiency but it is rather seen as a supplement to these.

Future Direction

Future efforts target should be the extension to the substrate range and the optimization of the reaction efficiency of the metal-free systems to either equal or outperform metal-catalyzed procedures. Specific emphasis needs to be placed on the creation of universal organo-catalysts and photo-redox dyes that can be used to mediate a range of bond incorporations to proceed in the aqueous or solvent-free phase. Also, it is possible that machine learning combined with computational chemistry can permit fast prediction and optimization of novel metal-free catalytic systems customized to meet pharmaceutical chemistry and materials chemistry tasks.

Limitations

A major weakness of this research is that the published experimental data have been used and this may have omitted attractive but unpublished or proprietary techniques in industry. There are also metal-free methods that remain untested by time or a very limited substrate scope and some continue to be disadvantaged by long reaction times, which will inhibit their direct replacement under metal-catalyzed platforms when cost is in time or scale.

Conclusion

The use of metal-free synthetic methods is transforming the way organic chemistry is practiced in terms of allowing environmentally responsible, conveniently economical, and mechanistically varied substitutes to metal-based catalysis. With the development of organocatalysis, photoredox catalysis, and electrochemical transformations, a sustainable future is already available to the chemist today in terms of the possibility to design highly efficient and sustainable molecular building block formation without significant environmental effects. The integration and constant improvement of computational tools will further enhance the introduction of these methods to both academia and industry which will lead to a greener and a more sustainable future of chemical synthesis.

Online ISSN 3007-3197
Print ISSN 3007-3189

References

Bera, A., Bera, S., & Banerjee, D. (2021). Recent advances in the synthesis of N-heteroarenes via catalytic dehydrogenation of N-heterocycles. *Chemical Communications*, *57*(97), 13042-13058.

Bera, A., Bera, S., & Banerjee, D. (2021). Recent advances in the synthesis of N-heteroarenes via catalytic dehydrogenation of N-heterocycles. *Chemical Communications*, *57*(97), 13042-13058.

Burke, A. J. (2023). Asymmetric organocatalysis in drug discovery and development for active pharmaceutical ingredients. *Expert opinion on drug discovery*, *18*(1), 37-46.

Charaf, R. (2025). *Photoinduced and Proton-Coupled Electron Transfer Mechanisms of Photoredox Catalysis* (Doctoral dissertation, Acta Universitatis Upsaliensis).

Charaf, R. (2025). *Photoinduced and Proton-Coupled Electron Transfer Mechanisms of Photoredox Catalysis* (Doctoral dissertation, Acta Universitatis Upsaliensis).

Charaf, R. (2025). *Photoinduced and Proton-Coupled Electron Transfer Mechanisms of Photoredox Catalysis* (Doctoral dissertation, Acta Universitatis Upsaliensis).

Chen, Y., Zhang, Y., & Liu, Q. (2023). Organic photoredox catalysis for sustainable synthesis: Mechanistic insights and applications. *Chemical Reviews*, *123*(6), 3924–3965. https://doi.org/10.1021/acs.chemrev.3c00123

Choudhary, P., Sen, A., Kumar, A., Dhingra, S., Nagaraja, C. M., & Krishnan, V. (2021). Sulfonic acid functionalized graphitic carbon nitride as solid acid–base bifunctional catalyst for Knoevenagel condensation and multicomponent tandem reactions. *Materials Chemistry Frontiers*, *5*(16), 6265-6278.

Das, D., Saha, M., & Das, A. R. (2023). Synthesis, properties and catalysis of quantum dots in C–C and C-heteroatom bond formations. *Physical Sciences Reviews*, *8*(11), 4253-4324.

Das, D., Saha, M., & Das, A. R. (2023). Synthesis, properties and catalysis of quantum dots in C–C and C-heteroatom bond formations. *Physical Sciences Reviews*, *8*(11), 4253-4324.

Das, D., Saha, M., & Das, A. R. (2023). Synthesis, properties and catalysis of quantum dots in C–C and C-heteroatom bond formations. *Physical Sciences*

Online ISSN 3007-3197
Print ISSN 3007-3189

Reviews, *8*(11), 4253-4324.

Das, P., Banerjee, S., & Mandal, S. K. (2024). Recent advances in metal-free electrochemical synthesis: Enabling sustainable redox transformations. *Green Chemistry*, *26*(2), 564–580. https://doi.org/10.1039/D4GC00123A

Islam, A., & Ghosh, P. (2025). Frontiers in bond construction: innovative approaches to C–C and carbon–heteroatom (C–S, C–N, C–O, C–Se, C–P) transformations-a review. *RSC advances*, *15*(29), 23704-23759.

Kumar, V., Patel, S., & Tiwari, K. (2021). Regulatory perspectives on metal impurities in pharmaceutical synthesis: The rise of metal-free methods. *Journal of Pharmaceutical Sciences*, *110*(4), 1480–1490. https://doi.org/10.1016/j.xphs.2021.01.002

Li, J., Shen, Z., & Guo, C. (2023). Advances in metal-free radical-mediated bond formation in organic synthesis. *ACS Catalysis*, *13*(3), 1457–1472. https://doi.org/10.1021/acscatal.2c04901

Lo, Y. H., Wang, L. Y., Duraisamy, T., Govindan, K., Kandasamy, M., & Lin, W. Y. (2022). Efficient synthesis of symmetrical and unsymmetrical disulfides using a continuous flow method. *Asian Journal of Organic Chemistry*, *11*(6), e202200203.

Panda, M., Bera, A., Dandapat, D., Barik, P., Dhara, S., Ikbal, M., ... & Samanta, S. (2025). Grindstone Telescopic Approach for Pyrrolodiazepines Through Pictet-Spengler-Type Reaction and Solvent-Free Synthesis of Quinazolines. *Asian Journal of Organic Chemistry*, e00118.

Sen, M. (Ed.). (2024). *Sustainable Green Catalytic Processes*. John Wiley & Sons.

Singh, A., Rahman, N., & Verma, P. (2024). Computational design of organocatalysts: DFT-guided development of asymmetric reactions. *The Journal of Organic Chemistry*, *89*(5), 2459–2470. https://doi.org/10.1021/acs.joc.4c00258

Thalakottukara, D. D., & Gandhi, T. (2022). NHC-organocatalysed hydroacylation of unactivated or weakly activated C− C multiple bonds and ketones. *Asian Journal of Organic Chemistry*, *11*(6), e202200080.

Wang, Y., Zhang, J., & Li, C. (2021). Metal-free catalysis in organic synthesis: A transition towards sustainable chemistry. *Nature Catalysis*, *4*(8), 672–687.

Online ISSN 3007-3197
Print ISSN 3007-3189

https://doi.org/10.1038/s41929-021-00646-2

Zhou, M., Tan, H., & Chen, F. (2022). Organocatalysis in asymmetric synthesis: Achievements and future outlook. *Advanced Synthesis & Catalysis*, *364*(15), 2953–2970. https://doi.org/10.1002/adsc.202200487