Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 9 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

# Advancements in Prenatal Development: Fetal Growth, Maternal Health, and Early Life Outcomes

**Rihana Dilshad**
Assistant Medical Superintendent, Sheikh Zayed Medical College, Rahim Yar Khan, Pakistan
**Sabira Sultana**
Department of Eastern Medicine, Faculty of Medical Sciences Government College University Faisalabad
**Amrat Sareen**
Department of Biology, HITEC School and Colleges, Taxilla Cantt, Rawalpindi, Pakistan
**Neelum Shahzadi**
Hamdrad University Islamabad Campus, Islamabad Pakistan
**Alia Hyder**
Hamdard University Islamabad Campus, Islamabad Pakistan
**Rizwana Dilshad**
Associate Professor, Swedish College of Pharmacy and Allied Health Sciences, Rahim Yar Khan, Pakistan
**Abdul Wadood**
Department of Eastern Medicine, Superior University Lahore, Pakistan
**Syed Muhammad Ali Shah**
Department of Eastern Medicine, Faculty of Medical Sciences Government College University Faisalabad
**Hamad Ahmed**
Department of Eastern Medicine, Faculty of Medical Sciences Government College University Faisalabad
**Muhammad Mudassar**
College of Allied Health Professionals, Faculty of Medical Sciences, Government College University Faisalabad, Pakistan
**Mudassar Khan**
Health Services Academy, Islamabad Pakistan
**Fahad Said Khan**
Department of Eastern Medicine, Faculty of Medical and Health Sciences, University of Poonch Rawalakot Azad Kashmir Pakistan
**Fethi Ahmet Ozdemir**
Department of Molecular Biology and Genetics, Faculty of Science and Art, Bingol University, Bingol, 1200, Türkiye
**Gaweł Sołowski**
Department of Molecular Biology and Genetics, Faculty of Science and Art, Bingol University, Bingol, 1200, Türkiye
**Muhammad Akram**
Department of Eastern Medicine, Government College University Faisalabad Pakistan
Corresponding Author: muhammadakram@gcuf.edu.pk

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 9 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189

http://amresearchreview.com/index.php/Journal/about

## ABSTRACT

The prenatal period represents a fundamental phase characterized by accelerated developmental transformations spanning from fertilization through parturition. This analysis explores the trio of distinct prenatal phases—germinal, embryonic, and fetal periods—emphasizing the significant morphological and physiological modifications occurring within each developmental window. The investigation further addresses the influence of various external determinants on fetal maturation, encompassing maternal psychological stress, teratogenic substance exposure, infectious agents, environmental contaminants, and nutritional status. The examination also incorporates the developmental origins of health and disease hypothesis, highlighting how early gestational disruptions may predispose individuals to subsequent pathological conditions, including neurodevelopmental disorders, cardiac abnormalities, and endocrine dysfunction. Comprehending these prenatal determinants is essential for enhancing perinatal outcomes and informing future research into the developmental programming of lifelong health trajectories.

**Keywords:** Prenatal development, germinal stage, embryonic stage, fetal stage, maternal health, environmental factors, fetal growth, birth defects, teratogens, maternal nutrition

## Introduction

The prenatal period, extending from conception to birth, typically spans around nine months and is marked by rapid and complex developmental processes. Growth during this phase is extraordinary: if the rate of prenatal growth observed in early gestation were sustained through the first year of life, an average newborn would weigh approximately 90 kg by their first birthday (Meredith et al., 1939). Fortunately, growth decelerates substantially in the final weeks before delivery..

Prenatal care represents a cornerstone of maternal healthcare systems, fundamentally designed to optimise maternal and fetal health outcomes throughout

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 9 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

gestation. Contemporary antenatal services encompass comprehensive surveillance for pregnancy-related complications, evidence-based interventions to mitigate risk factors, and structured educational programmes that enhance maternal preparedness for childbirth and parenting (Mohamed et al., 2025). The integration of digital health technologies into prenatal care delivery has transformed traditional service models, facilitating enhanced provider-patient communication, individualised care pathways, and real-time access to clinical decision-support tools (Field et al., 2025). This technological evolution has strengthened the capacity for shared decision-making between clinicians and pregnant women, ultimately contributing to improved perinatal outcomes and maternal satisfaction with care.

**The Course of Prenatal Development**

The prenatal period is typically divided into three developmental phases. The germinal stage spans the first two weeks following conception (1), the embryonic stage (2) extends from the second week to the end of the second month, and the fetal stage (3)continues from the beginning of the third month until birth. Key developmental milestones characterise each phase, as outlined below..

**Germinal Stage**

The first two weeks following conception are known as the germinal stage, which is the first stage of foetal development. When fertilization produces a zygote, this short stage starts. Rapid cell division starts in 36 hours. The zygote develops into a small mass of cells that proliferate (Mazia et al., 1961). This mass gradually moves into the uterus through the mother's fallopian tube. The cell mass starts to implant itself in the uterine wall on day six. This is not an automated procedure and takes around a week. Fewer than half of zygotes implant successfully. As a result, many pregnancies terminate before the women are aware that they have conceived.

Placental development initiates during the implantation process. This specialized organ facilitates the exchange of essential substances, including nutrients and oxygen, from maternal circulation to the developing fetus, while simultaneously

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 9 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

enabling the removal of metabolic waste products back to the mother's system. This critical biological exchange occurs through selective membrane barriers that maintain separation between fetal and maternal blood supplies, allowing molecular transfer while preventing the passage of cellular components.Gupta et al., 2020).

**Embryonic Stage**

The second stage of fetal development, known as the embryonic level, takes two weeks to end the second month. At this point, the developing organism, now known as embryo, begins to create most of its essential organs and physical systems. As cell division becomes increasingly specialised, structures like the brain, spine, and heart progressively come into being (Purves et al., 1988). Despite measuring only approximately one inch in length, the embryo displays recognizable human characteristics during this developmental phase. Limb buds have differentiated into distinct appendages, with digits becoming visible on both upper and lower extremities. Facial features, including ocular and auditory structures, are also discernible. This period represents a particularly vulnerable phase of development due to the concurrent establishment of essential organ systems and anatomical frameworks.The consequences can be disastrous if something disrupts normal development during the embryonic stage. This is when the majority of miscarriages happen. The majority of significant structural birth abnormalities are also caused by issues that arise during the embryonic period (Moore et al., 2016).

**Fetal Stage**

The third stage of pregnancy, known as the foetal stage, lasts for two months before delivery. Below is a summary of some salient features of foetal development. During the initial two months of bones and muscles build following embryonic development, the fetus undergoes rapid physical maturation. The skeletal framework begins to harden, enabling the developing organism to achieve mobility (Shea et al., 2015). Organ systems established in the preceding embryonic period continue their expansion and gradually become functional. Auditory capabilities, for example, emerge between the twentieth and twenty-fourth weeks of gestation.

The final trimester witnesses peak neural cell proliferation. During this period, the

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 9 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

pulmonary system matures while the central nervous system assumes greater regulatory control over bodily functions. Additionally, subcutaneous adipose tissue accumulates to provide thermal regulation. Somewhere between weeks 23 and 25, the fetus attains viability—the developmental milestone at which survival outside the womb becomes possible despite premature birth.

. The chance of survival is still low at 23 weeks (about 24%), but it rises rapidly to roughly 72% at 25 weeks. Sadly, many premature babies who are born around the viability barrier experience a variety of developmental issues (Gross et al., 1997). As an illustration, 43% of infants born between 20 and 24 weeks have neurodevelopmental deficiencies, while 19% have sensory deficiencies between 18 and 22 months.

**Environmental Factors and Prenatal Development**

The placenta connects the developing organism to its mother;events in the external world might have an indirect impact on the foetus, even if it develops inside the protective uterus.

**Nutrition**The growing foetus requires many vital nutrients, which makes maternal nutrition crucial. Severe maternal malnutrition raises the newborn's risk of neurological impairments and delivery difficulties (Chattopadhyay et al., 2016). Because moderate malnutrition is frequently confused with other risk factors linked to poverty, like drug usage and restricted access to healthcare, its effects are more challenging to quantify. Pregnant women should still eat a balanced diet that contains the necessary vitamins and minerals, even if they have easy access to food.

**Stress and Emotion**

Prenatal development may be impacted by maternal emotions in response to stressful situations, according to research. . Elevated maternal stress exposure during gestation correlates with increased fetal mortality rates, compromised immune system development, enhanced infectious disease susceptibility, delayed motor milestone acquisition, and impaired neurocognitive development (Goldenberg et al., 2005). The pathophysiological mechanisms underlying stress-induced prenatal complications involve disruption of maternal-fetal hormonal equilibrium, wherein maternal psychological distress triggers neuroendocrine cascade alterations that compromise optimal intrauterine developmental conditions.Supportive friendships

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 9 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

during pregnancy—through encouragement of healthy behaviours such as balanced eating and provision of emotional care—can enhance maternal mood and indirectly stabilise appetite. Conversely, limited social support from peers may worsen low mood and disrupt appetite regulation. Within the broader framework of social determinants of health, restlessness has been shown to correlate positively with friendship networks but negatively with educational attainment. Pregnant women with lower levels of education, often lacking access to reliable knowledge about pregnancy and childbirth, may struggle to cope effectively with the physical and psychological changes of pregnancy, thereby intensifying anxiety (Meng et al., 2025).

**Drug Use**

Maternal drug usage is another significant cause for worry. Sadly, the majority of medications taken by pregnant women can pass through the placental membranes. Almost all "recreational" substances have the potential to be harmful, but methamphetamine, cocaine, opioids, and sedatives are particularly risky (Bersani et al.., 2018). Over 21,000 newborns with narcotic addictions are born in the US each year as a result of maternal drug misuse. Additionally, this figure seems to be increasing quickly due to the opioid epidemic in the United States. Pharmacological exposures during pregnancy, including over-the-counter medications and prescription drugs administered for legitimate therapeutic indications, may present teratogenic risks requiring careful benefit-risk assessment. Maternal tobacco use demonstrates significant associations with adverse pregnancy outcomes, including increased rates of spontaneous abortion, fetal demise, preterm delivery, and sudden infant death syndrome. Long-term developmental consequences of prenatal tobacco exposure encompass compromised self-regulatory capacity, impaired academic achievement, and elevated risk for psychiatric disorders (Robson et al., 2020). However, establishing definitive causal relationships remains methodologically challenging due to frequent clustering of maternal smoking with additional high-risk behaviors and genetic susceptibility factors that confound observational analyses.Ndeke et al. (2024) identified carcinogens in electronic nicotine delivery system (ENDS) aerosols and marijuana vapour that are capable of crossing the placental barrier, thereby posing potential long-term cancer risks in offspring..

**Alcohol Consumption**

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 9 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

There are significant dangers associated with alcohol usage during pregnancy. The detrimental effects of high maternal alcohol use on the foetus have long been recognized. A collection of congenital (inborn) issues linked to heavy alcohol consumption during pregnancy is known as foetal alcohol syndrome (Abel et al., 2012). Microcephaly (small head), heart defects, irritability, hyperactivity, delayed motor development, and impaired cognitive development with academic challenges are common paediatric issues. Foetal alcohol syndrome may be far more common than previously believed, according to recent studies. During pregnancy, even modest alcohol consumption can have significant and long-lasting adverse effects. Moderate alcohol use during pregnancy has been linked to a higher risk of attention span, motor skill, and IQimpairments as well as an increase in impulsive, antisocial, and delinquent behaviour, according to research on the topic (Lynam et al., 2001).

**Maternal Illness**

Many, but not all, infectious pathogens are eliminated by the placenta. Thus, a variety of maternal illnesses may affect the development of the foetus. Measles, rubella, syphilis, and chickenpox are among the illnesses that can be harmful to the foetus; the type of harm varies depending in part on when the mother gets the illness (George et al., 2022). Pregnant women can also pass on the HIV that causes AIDS to their unborn children. AIDS can spread during pregnancy through the placenta, during childbirth, or through nursing.

Mother-to-child HIV transmission rates exemplify successful public health interventions in prenatal care, with vertical transmission declining from 20-30% in HIV-positive pregnant women during the pre-antiretroviral era to less than 1% in contemporary developed nations through implementation of highly active antiretroviral therapy during pregnancy and enhanced perinatal management protocols.

(Sebitloane et al., 2017).Chagas diseaseinfected by the mother without proper blood transfusion, due to positive antibodies *Trypanosoma cruzi*,and without were delivering babies with cardiac disease and Chagas diseases in 83% cases (Alavian et al., 2025).

**Environmental Toxins**

A growing body of evidence demonstrates that a wide range of environmental

Annual Methodological Archive Research Review
http://amresearchreview.com/index.php/Journal/about
Volume 3, Issue 9 (2025)

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

pollutants are detectable in utero, with significant implications for fetal and child development. Prenatal exposure to air pollution has been associated with impaired social competence and self-regulation in middle childhood, reduced cognitive performance at age five, and increased risk of obesity by age seven (Henrichs et al., 2015). Similarly, exposure to phthalates—commonly found in household products—has been linked to reductions in intelligence quotient, with reported losses of 6–7 points by age seven..The preventive role of nutrition in chronic diseases such as diabetes, hypertension, and cancer is well established. Increasing evidence also highlights its impact on embryonic brain development and the influence of both prenatal and preconceptional nutrition on susceptibility to disease. These effects extend beyond adequate caloric intake or maintaining a healthy body mass index, encompassing the quality and balance of macro- and micronutrient consumption, dietary patterns, and environmental exposures (Cernigliaro et al., 2025). Notably, factors such as heavy metals and volatile organic compounds, including butanone, may also contribute to long-term health outcomes (Wei et al., 2024).

**Fetal Origins of Adult Disease**

Most research on prenatal development has focused on its association with congenital anomalies and adverse outcomes manifesting in early life.
. The connections between prenatal variables and adult physical and mental health, however, are now being investigated by researchers. Research indicates that negative experiences during foetal development may "program" the foetal brain in ways that affect an individual's susceptibility to several diseases decades later (Buss et al., 2012). Epidemiological investigations demonstrate specific associations between intrauterine nutritional deficiencies and late-onset psychiatric disorders, including schizophrenia, which typically emerges during late adolescence or early adulthood. Low birth weight serves as a clinical indicator of multiple adverse prenatal exposures and correlates with elevated cardiovascular disease risk in later decades. Longitudinal cohort studies have established significant associations between disrupted fetal developmental processes and increased adult susceptibility to obesity, type 2 diabetes mellitus, major depressive disorder, bipolar affective

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

disorder, and specific malignancy subtypes.Faa et al. (2024) interpreted the effects of Thymosin beta-4 (TB4) as supportive evidence for the Barker hypothesis on the fetal origins of adult disease and its extension in the Developmental Origins of Health and Disease (DOHaD) theory. These findings suggest a potential shift in preventive strategies for major conditions such as atherosclerosis and its systemic complications affecting the heart, brain, and kidneys. It is increasingly evident that the prevention of many adult-onset diseases should begin in utero.Yaremenko et al. (2024) have linked abnormalities in fetal lung development to several chronic pulmonary conditions; for example, congenital diaphragmatic hernia (CDH) has been associated with pulmonary hypertension and adverse ventilatory outcomes.

(Capra et al., 2013). These discoveries on the foetal origin of disorders have spurred a reevaluation of the variables influencing health and illness. However, detractors note that research relating prenatal stress to adult health outcomes has mostly overlooked genetic susceptibilities to several disorders. The issue of causality is somewhat hazy because differences in how pregnant women respond to stress may be linked to their genetic susceptibilities to certain illnesses, which can be passed down to their children.Establishing causal relationships between prenatal exposures and adult health outcomes presents methodological challenges due to prolonged follow-up requirements, limited historical data availability, and complex confounding variables inherent in longitudinal research designs. Prospective cohort studies necessitate recruitment of substantial sample sizes (for example group of 100 foetus until their death) and comprehensive lifelong surveillance protocols to achieve statistical power for detecting long-term associations (Bilinski et al., 2025). Such investigations encounter significant ethical considerations regarding informed consent, particularly when participants transition from childhood to autonomous adulthood, requiring ongoing consent processes and adherence to evolving research ethics standards throughout extended observation periods.

**Conclusion**

Prenatal development represents a critical developmental window characterized by accelerated cellular proliferation, organogenesis, and morphological differentiation that establishes the physiological framework for lifelong health trajectories. The sequential progression through germinal, embryonic, and fetal developmental phases

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

demonstrates distinct vulnerability periods wherein intrinsic biological programming intersects with extrinsic environmental influences. While endogenous genetic and epigenetic mechanisms orchestrate fundamental developmental processes, maternal environmental exposures including nutritional status, psychosocial stressors, teratogenic substances, infectious agents, and environmental toxicants exert profound effects on fetal neurodevelopment and organ system maturation. Adverse intrauterine conditions demonstrate persistent consequences extending beyond the neonatal period, manifesting as altered neurocognitive performance, compromised physical development, and increased susceptibility to psychiatric disorders throughout the lifespan. The developmental origins of health and disease hypothesis provides compelling evidence linking prenatal environmental perturbations to increased risk for cardiovascular disease, metabolic disorders, and neuropsychiatric conditions in adulthood. Contemporary research emphasizing fetal programming mechanisms highlights the critical importance of comprehensive antenatal care and maternal health optimization in primary prevention strategies for chronic disease burden. Understanding these complex gene-environment interactions during prenatal development informs evidence-based interventions targeting maternal healthcare delivery, environmental risk reduction, and population health approaches designed to enhance developmental outcomes across generations..

**Reference**

Abel EL. Fetal alcohol syndrome and fetal alcohol effects. Springer Science & Business Media; 2012 Dec 6.

Alavian, N., Izatt, S., Rolfe, R. J., Hernandez, J., Kurtzberg, J., Martin, L., & Livingston, E. (2025). Chagas disease screening of maternal donors in publicly banked umbilical cord blood in North Carolina, United States 2007–2023. *Pregnancy*, *1*(4), e70045.

Bersani FS, Imperatori C. Misuse, recreational use, and addiction in relation to prescription medicines. In Handbookof Novel Psychoactive Substances 2018 Oct 10 (pp. 364-381). Routledge.

Bilinski, A., Emanuel, N., & Ciaranello, A. (2025). Sins of Omission: Model-Based Estimates of the Health Effects of Excluding Pregnant Participants From Randomized Controlled Trials. *Annals of Internal Medicine*, *178*(6), 868-877.

Buss C, Entringer S, Wadhwa PD. Fetal programming of brain development: intrauterine stress and susceptibility to psychopathology. Science signaling. 2012 Oct 9;5(245):pt7-.

Capra L, Tezza G, Mazzei F, Boner AL. The origins of health and disease: the

Online ISSN 3007-3197
Print ISSN 3007-3189
http://amresearchreview.com/index.php/Journal/about

influence of maternal diseases and lifestyle during gestation. Italian journal of pediatrics. 2013 Dec;39:1-2.

Cernigliaro, F., Santangelo, A., Nardello, R., Lo Cascio, S., D'Agostino, S., Correnti, E., Marchese, F., Pitino, R., Valdese, S., Rizzo, C., Raieli, V., & Santangelo, G. (2024). Prenatal Nutritional Factors and Neurodevelopmental Disorders: A Narrative Review. *Life*, *14*(9), 1084. https://doi.org/10.3390/life14091084

Chattopadhyay N, Saumitra M. Developmental outcome in children with malnutrition. Journal of Nepal Paediatric Society. 2016 Dec 31;36(2):170-7.

Faa, G., Fanos, V., Manchia, M., Van Eyken, P., Suri, J. S., & Saba, L. (2024). The fascinating theory of fetal programming of adult diseases: A review of the fundamentals of the Barker hypothesis. *Journal of Public Health Research*, *13*(1), 22799036241226817.

Field, N. K., Venkatesan, C., Gano, D., Agarwal, S., Young, K. A., Wheeler, S., ... & Lemmon, M. E. (2025). Communicating neurological prognosis in the prenatal period: a narrative review and practice guidelines. *Pediatric Research*, 1-17.

George CR, Jeffery HE, Lahra MM. Infection of mother and baby. Keeling's Fetal and Neonatal Pathology. 2022:207-

Goldenberg RL, Culhane JF, Johnson DC. Maternal infection and adverse fetal and neonatal outcomes. Clinics in perinatology. 2005 Sep 1;32(3):523-59.

Gross RT, Spiker D, Haynes CW. Helping low birth weight, premature babies: The Infant Health and Development Program. Stanford University Press; 1997.

45.

Gupta PD. What Your Blood Tells? A Review. Journal of Cell and Tissue Research. 2020;20(2):6897-913.

Henrichs J, Van den Bergh BR. Perinatal developmental origins of self-regulation. Handbook of biobehavioral approaches to self-regulation. 2015:349-70.

Lynam DR, Henry B. The role of neuropsychological deficits in conduct disorders. Conduct disorders in childhood and adolescence. 2001:235-63.

Mazia D. Mitosis and the physiology of cell division. InThe cell 1961 Jan 1 (pp. 77-412). Academic Press.

Meng, W., Shalayiding, S., Wang, X. *et al.* Relationship between prenatal anxiety, depression, pregnancy stress and their social determinants. *BMC Psychol* **13**, 905 (2025). https://doi.org/10.1186/s40359-025-03237-3

Meredith HV, Brown AW. Growth in body weight during the first ten days of postnatal life. Human Biology. 1939 Feb 1;11(1):24.

Mohamed, H., Ismail, A., Sutan, R. *et al.* A scoping review of digital technologies in antenatal care: recent progress and applications of digital technologies. *BMC Pregnancy Childbirth* **25**, 153 (2025). https://doi.org/10.1186/s12884-025-07209-8

Moore KL. Before We Are Born: Essentials of Embryology and Birth Defects: First

Online ISSN 3007-3197
Print ISSN 3007-3189



South Asia Edition E-Book. Elsevier Health Sciences; 2016 Mar 23.

Ndeke, J. M., Klaunig, J. E., & Commodore, S. (2024). Nicotine or marijuana vaping exposure during pregnancy and altered immune responses in offspring. *Journal of environmental exposure assessment*, *3*(1), 10-20517.

Purves D. Body and brain: a trophic theory of neural connections. Harvard University Press; 1988.

Robson DA, Allen MS, Howard SJ. Self-regulation in childhood as a predictor of future outcomes: A meta-analytic review. Psychological Bulletin. 2020 Apr;146(4):324.

Sebitloane HM, Moodley D. The impact of highly active antiretroviral therapy on obstetric conditions: a review. European Journal of Obstetrics & Gynecology and Reproductive Biology. 2017 Mar 1;210:126-31.

Shea CA, Rolfe RA, Murphy P. The importance of foetal movement for co-ordinated cartilage and bone development in utero: clinical consequences and potential for therapy. Bone & joint research. 2015 Jul 1;4(7):105-16.

Wei, S., Li, Z., Murugappan, K., Li, Z., Lysevych, M., Vora, K., ... & Fu, L. (2024). Nanowire array breath acetone sensor for diabetes monitoring. *Advanced Science*, *11*(19), 2309481.

Yaremenko, A.V.; Pechnikova, N.A.; Porpodis, K.; Damdoumis, S.; Aggeli, A.; Theodora, P.; Domvri, K. Association of Fetal Lung Development Disorders with Adult Diseases: A Comprehensive Review. *J. Pers. Med.* **2024**, *14*, 368. https://doi.org/10.3390/jpm14040368